



British Association of
Perinatal Medicine

Perinatal Management of Extreme Preterm Birth Before 27 Weeks of Gestation

A BAPM Framework for Practice

October 2019
Updated August 2025

Contents

Foreword for revised 2025 Framework	3
Executive summary	4
Members of the 2025 update Working Group (in alphabetic order)	6
Members of the 2019 Working Group.....	7
Introduction	8
Risk-based approach to decision-making	11
Obstetric management	18
Neonatal management	21
Implications of this framework for Transport/LNU and SCU teams	24
Implementation of this Framework for Practice	24
Recommendations for audit and research	25
Appendix 1: Outcomes for extremely preterm babies: international comparisons.....	26
Appendix 2: Example case scenarios	34
Appendix 3: Situations of uncertainty and potential conflict: recommended considerations and actions.....	37
Appendix 4: Communication: Guidance for professionals consulting with families at risk of extreme preterm birth.	40
Appendix 5: Helping parents to understand extreme preterm birth.	43
References	48

Developed in Partnership with: (To be confirmed)

Endorsed by: (To be confirmed)

Foreword for revised 2025 Framework

The 2019 BAPM Framework on Perinatal Management of Extreme Preterm Birth before 27 weeks gestation provided a revised approach to decision-making and care for some of the smallest and most vulnerable premature babies. It was based on evidence from UK and international data on improved survival, and included consideration of survival focused care for babies as early as 22 weeks gestation, following a risk-stratified approach to counselling and decision-making. The change in guidance had wider implications beyond neonatal services, with an increase in antenatal transfers of women at 22-23 weeks gestation with threatened preterm labour. We recommend that perinatal teams across neonatal and maternity services familiarise themselves with this framework to facilitate counselling and management decisions.

Whilst the risk stratified approach was welcomed by perinatal teams, feedback indicated that this needed greater emphasis and clarity to avoid misinterpretation as a recommendation for universal survival focused care. Whilst this update retains the core elements of the 2019 Framework, we have made revisions to provide clearer language, support and guidance around appropriate decisions for babies assessed to be at extremely high risk of dying. It emphasises that for such babies, comfort care should be provided. This framework also includes updated survival and neurodevelopmental outcome figures and a revised infographic including information on the proportion of babies liveborn. This revision emphasises the importance of joint maternity and neonatal counselling with families, which includes communication between centres when transfers antenatally or postnatally are considered.

To help support teams apply the framework in practice, we have included new example cases ([Appendix 2](#)), and situations that present uncertainty and potential conflict, for example born outside hospital or born at 22 weeks in centres without a co-located NICU ([Appendix 3](#)). We have also included signposts to available educational and training resources.

Neonatology continues to evolve as a specialty, and it is inevitable that this framework will need further updates as new evidence becomes available. The key elements, however, will remain constant: the need for perinatal teams to work collaboratively, to inform and involve parents in decision-making, and to support them at one of the most stressful times of their lives.

Executive summary

1. Decision making for babies born before 27 weeks of gestation should not be based on gestational age alone, but on assessment of the baby's prognosis taking into account multiple factors.
2. Neonatal stabilisation may be considered for babies born from 22 weeks of gestation following assessment of risk and multiprofessional discussion with parents.
3. While the main focus of this framework is on the care of babies once they arrive to hospital or that are born in hospital, some scenarios of babies born outside a hospital setting are covered in [Appendix 3](#).
4. Decisions should be made with input from obstetric and neonatal teams, and joint counselling with families is recommended. Where transfers antenatally or postnatally are considered, communication between referring, receiving and transfer centres should take place to facilitate shared and informed decision making with families.
5. Risk assessment should be performed with the aim of stratifying into three groups:
 - a) Extremely high risk.
 - b) High risk.
 - c) Moderate risk.
6. For babies at **extremely high risk** of poor outcome, comfort focused care should be provided. Survival focused care should not be offered. This includes all babies born before 22 weeks. It also includes some babies born at 22 weeks or later with extremely low chance of survival without severe impairment. A full assessment of the risks of continuation of the pregnancy to the mother's health must be considered alongside the outcome for the baby, with particular attention to preterm prelabour rupture of membranes (PPROM) and the development of infection.
7. For babies at **high risk** of poor outcome, the decision to provide either survival focused care or comfort focused care should be based on individual assessment and informed by parental wishes, and supported by consultation and in partnership with obstetric and neonatal professionals.
8. For babies at **moderate risk**, survival focused care should be provided.
9. If survival focused care for the baby is anticipated, pregnancy and birth should be managed with the aim of optimising the baby's condition at birth and subsequently.
10. If comfort focused care for the baby is planned, obstetric management should focus on maternal wellbeing and minimising risks for future pregnancies.
11. Whenever possible, babies born before 27 weeks (where survival focused care is planned) should be born in a maternity facility co-located with a designated neonatal intensive care unit (NICU).
12. A new section on Implications for Transport/LNU and SCU teams has been included.
13. Conversations with parents should be clearly documented and care taken to ensure that the

agreed management plan is communicated between professionals and staff shifts.

14. Management should be regularly reviewed before and after birth in conjunction with the parents; plans may be reconsidered if the risk for the baby changes, or if parental wishes change.
15. Perinatal professionals involved in the counselling and care of families and babies born before 27 weeks should undertake relevant training and engage in educational opportunities.
16. References have been updated and include signposting to previous BAPM frameworks and other national guidance.
17. Recommendations for research have been included in line with all BAPM frameworks.

DRAFT

Members of the 2025 update Working Group (in alphabetic order)

Dr David Bartle

Consultant Paediatrician & Neonatologist, Royal Devon University Healthcare NHS Foundation Trust, BAPM LNU & SCU Representative

***Dr Sarah Bates**

Consultant Paediatrician and Neonatologist, Great Western Hospital, Swindon, BAPM Perinatal Optimisation Lead for Quality Team

Dr Cheryl Battersby

Consultant Neonatologist, Chelsea & Westminster, NIHR Clinician Scientist, Clinical Associate Professor, Imperial College London, BAPM Honorary Secretary and Chair of the BAPM Nanopreterm Specialist Interest Group

Dr Allan Jackson

Consultant Neonatologist, Princess Royal Maternity Hospital, Glasgow, Neonatal Clinical Lead, ScotSTAR Transport Service

Dr Tracey Kay

Consultant Obstetrician, Royal Devon University Healthcare NHS Foundation Trust, Obstetric Representative to BAPM Quality Team

***Ms Kate Mulley**

Director of Research, Education and Policy for Sands

Dr Sam Oddie

Consultant Neonatologist, Bradford Royal Infirmary, Clinical Lead for National Neonatal Audit Project, BAPM Data Lead

***Professor Lucy K Smith**

Professor of Perinatal Health, Department of Population Health Sciences, University of Leicester representing MBRRACE-UK

Dr Steve Wardle

Consultant Neonatologist, Nottingham University Hospitals, BAPM President Elect

***Professor Dominic Wilkinson**

Professor of Medical Ethics, University of Oxford and Consultant Neonatologist, John Radcliffe Hospital, Oxford

*Involved in both 2019 and 2025 frameworks.

Members of the 2019 Working Group

Ms Caroline Lee-Davey

Chief Executive Officer of Bliss

Ms Erica Everett

Neonatal Nurse, East of England Operational Delivery Network, representing the Neonatal Nurses Association

Dr Tracey Johnston

Consultant in Maternal and Fetal Medicine, Birmingham Women's and Children's NHS Foundation Trust, representing British Maternal and Fetal Medicine Society

Dr Helen Mactier (Chair)

Consultant Neonatologist and Honorary Clinical Associate Professor, Princess Royal Maternity and the University of Glasgow. President of BAPM

Prof Neil Marlow

Professor of Neonatal Medicine, University College London, representing RCPCH

Dr Tara Selman

Consultant in Fetal and Maternal Medicine University Hospital Southampton, representing British Maternal and Fetal Medicine Society

Dr Meekai To

Consultant in Obstetrics and sub specialist in fetal and maternal medicine, King's College Hospital, London, representing the Royal College of Obstetricians and Gynaecologists

The working group wishes to acknowledge extremely valuable contributions from Dr Emily van Blankenstein and Dr Helen McDermott in developing the evidence synthesis in [Appendix 1](#) and for formatting the references. We would like to acknowledge the invaluable, voluntary assistance of Phil Johns, Graphic Designer, Managing Director of Respond in preparing the infographics. We are also grateful to Caroline Lee-Davey (BLISS) for reviewing and contributing to the revision, and to Kate Dinwiddy, Chief Executive of BAPM, and the BAPM office team for administrative support.

Introduction

The BAPM 2019 Framework for Practice was developed by consensus, with wide consultation, taking into account available outcome data at the time (2016 births) both from the UK and internationally. We are grateful for input from the RCOG, the British Maternal and Fetal Medicine Society, the Royal College of Paediatrics and Child Health, MBRRACE-UK, UK Pre-Hospital Maternity and Newborn Care Group, the Neonatal Nurses Association and parent representative organisations including Bliss and Sands. The scope includes births up to 26⁺⁶ weeks of gestation, better to align with national recommendations and published data (Neonatal Critical Care Review, 2019).

Care of the baby, woman and family around the time of an extremely preterm birth is one of the most challenging aspects of perinatal medicine, both for clinicians and families. In 2006, the Nuffield Council on Bioethics convened a working group to explore the ethical, social, economic and legal issues around clinical decisions made in fetal and neonatal medicine (Nuffield Bioethics 2006); in response to their report the British Association of Perinatal Medicine (BAPM), in conjunction with other professional groups, developed a Framework for Clinical Practice for the Management of Babies born Extremely Preterm at less than 26 weeks of gestation. These documents were based largely on data from the original EPICure study in 1995 of births before 26 weeks of gestation, with some additional preliminary data from EPICure 2 regarding babies born in 2006 before 27 weeks of gestation (Costeloe, 2000, Marlow, 2005).

The ethical principles that formed the basis for these earlier guidelines have not changed but advances in perinatal care have led to improved outcomes for babies admitted to UK neonatal intensive care units (NICUs), particularly at the lowest gestational ages. Reports from other countries confirm increased survival and improved neurodevelopmental outcome for babies born before 27 weeks of gestation (Norman, 2019, Patel, 2017, Mehler, 2016, Myrhaug, 2019). Although internationally there remain differences in practice, there is increasing willingness to consider stabilisation at birth and subsequent intensive care for the most extremely preterm babies (Guillén, 2015, Lemyre, 2017, Wilkinson, 2018), accompanied by greater acknowledgement of the importance of involving parents in perinatal decision making (POPPY Steering Group 2018). Reported outcomes are, of course, impacted by willingness to consider survival focused interventions before and after birth (Rysavy, 2015).

The 2019 BAPM Perinatal Management of Extreme Preterm Birth Before 27 Weeks of Gestation Framework, updated its previous guidance from 2008, and recommended consideration of neonatal stabilisation of some babies from 22 weeks of gestation and above according to the results of risk stratification and following multi-professional discussion with parents. Recent publications combining UK data from Mothers and Babies: Reducing Risk through Audits and Confidential Enquiries (MBRRACE-UK) and the National Neonatal Research Database (NNRD) provide evidence of a three-fold increase in babies born at 22 weeks of gestation given survival focused care and admitted to neonatal units since the introduction of the BAPM 2019 framework (Smith, 2023). Two out of 10 babies provided with survival focused care and 3 out of 10 babies admitted to neonatal care survived to neonatal discharge. This corresponded to 13 and 39 babies in years 2018-19 and 2021, respectively. There has been no change over time in the provision of survival focused care for babies alive at the onset of labour born at 23 and 24 weeks (80% and 90%, respectively) (Smith, 2023). The BAPM 2019 framework had implications on the maternity service, with an increased need for capacity to transfer women antenatally between 22 and 23 weeks gestation with threatened preterm labour, to NICUs (Griffin, 2025).

In addition to the latest published figures (Smith, 2023) for babies born 22-24 weeks, this framework also includes updated survival figures for babies born in 2020-21 at 25 and 26 weeks in England and

Wales; revised infographics providing appropriate population denominators including proportion of babies liveborn, useful for counselling parents at different time points along the perinatal pathway; additional guidance for specific scenarios in the appendices. The framework emphasises and encourages perinatal teams to carefully consider risk factors (and not just gestational age) for decisions about survival focused care in the most vulnerable cohort of babies.

Prevention of preterm birth remains a national priority and all maternity services should ensure that measures are in place to realise this ambition. National guidance is available to enable prevention strategies; this guidance focuses on the importance of good communication between professionals and parents, strategies to ensure high quality survival focused and/or comfort focused care as appropriate and interventions to optimise outcomes for babies born too soon (NICE NG25, NHS England Saving Babies Lives Version 3).

Perinatal care at extremely preterm gestations will always need to be individualised and should be led by senior staff in midwifery, obstetrics and neonatology. Joint counselling with maternity/obstetric and neonatal teams to inform management decisions for mother and baby is recommended where possible. Parents should be included in discussions about perinatal care, and their hopes and expectations explored with honesty and compassion in a realistic way. Decisions should be made together with parents, based on the best available evidence about the prognosis for the individual baby, and mindful of the need to act in the baby's best interests.

Remit

The purpose of this Framework for Practice is to assist decision-making prior to and/or at the time of birth relating to perinatal care and preterm birth at 26 weeks and 6 days of gestation or less in the United Kingdom. Options for the pregnancy should have been discussed with the parents. Decision-making around termination of pregnancy is out of scope of this framework. For some extremely preterm babies, postnatal events may indicate that continuation of neonatal intensive care is not in the baby's best interests. While parents should be made aware of this possible outcome, this Framework does not address decisions around withdrawal or withholding of life-prolonging treatment after a baby has been admitted to a neonatal unit. A separate recent BAPM framework '[Recognising uncertainty: an integrated framework for palliative care in perinatal medicine](#)' outlines the principles of comfort focused care and support for these babies.

Whilst this framework focuses on the management of extremely preterm babies after arrival at hospital or those born in hospital, [Appendix 3](#) also includes discussion of the care of extremely preterm babies who have been transferred after being born out of hospital. Management of extremely preterm babies by pre-hospital clinicians is found in the BAPM Framework: '[Pre-hospital management of the baby born at extreme preterm gestation](#)'.

We have included guidance designed to assist health professionals in communicating with parents about the issues and information contained within this document. It is emphasised that each case will be unique, and that communication should always be tailored accordingly. We hope that this 2025 update to the 2019 framework, will be incorporated into local and network guidelines, to ensure consistency of practice within units and networks and acknowledgement of the importance of individualised care for families.

Definitions

In the UK, a **stillbirth** is legally defined as the birth of a baby with no signs of life at, or after, 24 completed weeks of pregnancy.

Within the document “**parents**” refers to the mother or birthing person and their partner. BAPM is aware that the use of gendered language such as mother, can make some families feel excluded. When supporting individual families, professionals should use the terms that the family identifies with, as well as their desired pronouns.

We have used the terms “**survival focused care**” to refer to obstetric and neonatal management that has the aim of sustaining life for the baby, and “**comfort focused care**” to refer to obstetric and neonatal management when the aim is not to attempt to sustain the life of the baby, but to focus on the baby’s comfort.

“**NICU**” refers to a designated neonatal intensive care unit, sometimes termed a level 3 unit.

Perinatal care teams include multi-professional healthcare staff dedicated to providing care for pregnant women, their newborns, and their families. The team includes, but is not limited to, obstetricians, midwives, neonatologists/paediatricians, nurses, and allied health professionals including dietitians, physiotherapists, and occupational therapists. Importantly, parents are also considered integral members of the team, actively involved in decision-making and care planning.

Risk-based approach to decision-making

A key ethical consideration for decisions about instituting life-sustaining for an extremely preterm baby is the baby's prognosis – the likelihood of an acceptable (or unacceptable) outcome for baby and parents. If there is a plan to provide life-sustaining treatment for the baby, then it follows that the pregnancy and birth should be managed with the aim of optimising the baby's condition at birth and subsequently.

We advise a stepwise approach to decision-making, involving three key stages:

1. Assessment of the risk for the baby if birth occurs, incorporating both gestational age and factors affecting fetal and/or maternal health.
2. Counselling parents, and their involvement in decision-making. It is important that parents are offered choices and supported to make decisions appropriate for their individual preferences.
3. Agreeing and communicating a management plan.

1. Assessment of the risk for the baby

1a. Gestation-based risk assessment, including mortality and survival with severe impairment

Gestation

In accordance with NICE guidelines, all pregnant women in the UK should have been offered an early ultrasound scan between 10⁺⁰ and 13⁺⁶ weeks of gestation, with crown–rump length (CRL) measurement used to determine gestation. This assessment is accurate to within 5 days in 95% of cases (NICE NG201). From 14⁺⁰ weeks (CRL > 84mm), gestation should be estimated from fetal head circumference; the estimated uncertainty of this gestation prediction is 6-7 days at 14 weeks, rising to 12-14 days by 26 weeks of gestation (Papageorghiou, 2016).

Survival

Survival of extremely preterm babies has increased steadily since 2006 with greater willingness to offer neonatal intensive care. The number of surviving babies at 22 weeks of gestation remains small, with appreciable in-labour mortality. UK figures accord with international data ([Appendix 1](#)). Survival to live birth is influenced by management of labour and birth, and survival after birth will clearly be affected by whether active survival focused care is provided.

Management should be based on the most recent data available. Recent UK data for babies born 2020-2021 is included in [Table 1](#) in Appendix 1. Survival rates differ by denominator. For babies born at 22 weeks of gestation, survival to neonatal unit discharge is around 8% of babies alive at the onset of labour, 10% of live births, 20% of babies who received survival focused care, and 30% of babies admitted to neonatal units. Survival rates are higher for babies born at greater gestation ([Appendix 1](#)). As a proportion of neonatal admissions, survival is around 50% for babies born at 23 weeks, 70% at 24 weeks, 80% at 25 weeks and 85% at 26 weeks. Denominators used should be relevant to the decision being made. For decisions about whether to provide survival-focused care at birth, the most relevant statistic is the chance of survival (or severe morbidity) for live born babies who receive survival focused care.

Morbidity

1 in 10 of babies born at 22 weeks gestation and 1 in 5 of those born at 23 weeks of gestation admitted to neonatal care survive to discharge without major morbidities (excluding BPD). All

surviving babies at 22 and 23 weeks of gestation have evidence of BPD (Smith, 2023).

There are important differences in individual views about acceptable levels of disability. What for one individual or family may be an acceptable outcome may not be for another. For decisions about provision of potentially life-sustaining treatment, the ethically relevant consideration is the risk of disabilities that could affect whether it is in the baby's best interests to survive and thus risk assessment should focus on the most severe disabilities (Wilkinson, 2013).

In the absence of regularly updated national data on the prevalence of severe disability after extremely preterm birth, we recommend that the well-established "severe impairment" category, as defined by the 2008 BAPM Working Group, be used to inform parents when discussing risk following extremely preterm birth. It is acknowledged that many more extremely preterm babies will be affected by milder degrees of disability; this should also be included in information provided to parents, with clear explanation that disability is generally impossible to predict for individual babies at birth.

The severe impairment category includes any of:

- Severe cognitive impairment with an IQ lower than 55 (< -3 standard deviation); this will usually result in the need for special educational support and require supervision in daily activities.
- Severe cerebral palsy – classified as Gross Motor Function Classification System (GMFCS) grade 3 or greater ([Appendix 1](#)).
- Blindness or profound hearing impairment.

The risk of severe impairment increases with increasingly preterm birth ([Appendix 1](#)) and using the most recent population level two-year neurodevelopmental data from babies born 2008-2018 is approximately the following for babies who receive survival focused care and survive to two years of age (van Blankenstein, 2024).

22⁺⁰ - 22⁺⁶ weeks:	1-in-3 survivors has severe impairment*
23⁺⁰ - 23⁺⁶ weeks:	1-in-4 survivors has severe impairment
24⁺⁰ - 24⁺⁶ weeks:	1-in-5 survivors has severe impairment
25⁺⁰ - 25⁺⁶ weeks:	1-in-9 survivors has severe impairment.
26⁺⁰ - 26⁺⁶ weeks:	1-in-12 survivors has severe impairment.

*Presently, relatively few babies born at 22 weeks of gestation have available long-term outcome data available, meaning estimates of rates of such adverse outcome are imprecise. However, it is believed that the proportion of such babies with severe impairment is at least 30%.

1b. Modified risk assessment

Accurate information about the current pregnancy, including assessment of both fetal and maternal health should be used to refine gestation-based risk of absolute survival and survival without severe impairment.

A range of factors are associated with increased or decreased risk:

Fetal factors which may increase risk include male sex, multiple pregnancy, congenital anomaly and poor fetal growth.

Clinical conditions which pose additional risk and have been associated with increased mortality and

morbidity include the following:

Chorioamnionitis increases the risk of neonatal complications including mortality and morbidity, particularly brain injury and adverse long-term neurodevelopment outcomes. Although ultimately a histological diagnosis (hence not possible to diagnose prior to birth), combinations of clinical symptoms can lead to a clinical diagnosis of chorioamnionitis in the antenatal or intrapartum periods. Symptoms include lower abdominal pain or uterine tenderness, abnormal vaginal discharge, fever, malaise, reduced fetal movements, increased baseline fetal heart rate, abnormal maternal observations, abnormal maternal biochemistry (in particular, C-reactive protein and white cell count) (RCOG GTG No.73).

Preterm prelabour rupture of membranes (PPROM): Pregnancies with PPRM prior to 23 weeks gestation have a high risk of perinatal and maternal morbidities. Risk is highest with very early PPRM (eg <20 weeks) in association with oligo or anhydramnios (Goodfellow, 2024).

Therapeutic strategies: administration of antenatal steroid and magnesium sulphate are associated with improved survival and neonatal outcomes as well as reduced risk of childhood impairment, even before 24 weeks of gestation (Roberts, 2017, Travers, 2017, Ehret, 2018, Doyle, 2019).


Clinical Setting: survival is highest at these extreme preterm gestations in centres with experienced staff and higher patient numbers. A strategy of in utero transfer below 27 weeks of gestation for birth in a maternity unit with a co-located NICU is strongly recommended (where survival focused care is clinically appropriate and desired by parents Marlow, 2014, David, 2018, Helenius, 2019, BAPM NSQI 2017). Where extremely-preterm babies are born in centres without a co-located NICU, this will increase the risk for the babies, and in some cases will mean that survival focused care is not appropriate. [See Appendix 3.](#)

Following full history taking and risk assessment, the risk of unacceptably poor outcome if life-sustaining care is provided for the baby will generally fall into one of the following categories:

- Extremely high risk.
- High risk.
- Moderate risk.

A visual tool for refinement of risk is illustrated in [Figure 1](#). Some clinical factors (chorioamnionitis, PPRM) are not included in the visual tool because of variability in diagnosis and impact on risk. Nevertheless, it would be important to include these in assessment of risk in cases where they clearly adversely affect individualised risk.

Figure 1: Visual tool for refinement of risk



Box 1 represents the consensus of the Working Group in regard to risk categories for the purposes of this framework.

BOX 1

Extremely high risk: The Working Group considered that babies with a > 90% chance of either dying or surviving with severe impairment if liveborn and survival focused care is instituted would fit into this category. For example, this would include:

- All babies born at less than 22 weeks of gestation.
- Babies born at 22⁺⁰ - 22⁺⁶ weeks of gestation with unfavourable risk factors (for example, those born in hospitals without a NICU and/or in the first half of the week, and/or in the absence of antenatal steroids).
- Some babies born at 23⁺⁰ - 23⁺⁶ weeks of gestation with unfavourable risk factors, including severe fetal growth restriction.
- (Rarely) babies born at or after 24⁺⁰ weeks of gestation with significant unfavourable risk factors, including very severe fetal growth restriction.

High risk: The Working Group considered that babies with a 50-90% chance of either dying or surviving with severe impairment if survival focused care is instituted would fit into this category. For example, this would include:

- Babies at 22⁺⁰ - 23⁺⁶ weeks of gestation with favourable risk factors.
- Some babies \geq 24⁺⁰ weeks of gestation with unfavourable risk factors and/or co-morbidities.

Moderate risk: The Working Group considered that babies with a < 50% chance of either dying or surviving with severe impairment if survival focused care is instituted would fit into this category. For example, this would include:

- Most babies \geq 24⁺⁰ weeks of gestation.
- Some babies at 23⁺⁰ - 23⁺⁶ weeks of gestation with favourable risk factors.

For women presenting to a maternity centre without a co located NICU, assessment of risk should include early discussion with the relevant referral centre. For pregnancies from 22⁺⁰ weeks of gestation, decisions should not be based on gestational age alone. Within a multiple pregnancy, the risk may differ between fetuses and so each should be considered as an individual. This means that appropriate management may not be the same for each baby, even with the same gestational age. If birth occurs prior to 22⁺⁰ weeks of gestation, survival focused neonatal care is not appropriate.

The agreed risk for the baby has ethical and practical implications for the options that should be available.

Extremely high risk: For babies with an extremely high risk of death or of survival with unacceptably severe impairment despite treatment, comfort-focused care would be in the best interests of the baby. Survival focused care should not be offered. There is no absolute indication for paediatric/neonatal attendance at the birth although for individual families this may be helpful.

High risk: For babies with a > 50% risk of death or of surviving with unacceptably severe impairment despite treatment, it is uncertain whether survival focused care is in the best interests of the baby

and their family. Parents should be counselled carefully and parental wishes should inform a joint decision to provide either survival focused or comfort focused care. Ideally, a senior neonatal clinician who has previously met the parents will be available to attend the birth and supervise implementation of the agreed plan, together with an experienced neonatal team.

Moderate risk: For babies with a < 50% risk of death or of survival with unacceptably severe impairment, survival focused care would be in the best interests of the baby. A senior neonatal clinician should attend the birth, together with an experienced neonatal team.

2. Counselling parents and decision-making

Whenever possible, parents should be involved in planning an extremely preterm birth. The planning consultation should include senior clinical staff from the obstetric, midwifery and neonatal teams who will be caring for the mother and her baby before, during and after the birth.

The assessed category of risk to the baby (including the inherent uncertainty around this) should be conveyed sympathetically and with clarity, and the hopes and expectations of parents explored with honesty and compassion in a realistic way. Clear, balanced information should be shared and management options discussed. Time should be allowed for clarification and questions, and parents offered the opportunity to revisit discussions with the perinatal team at any point, acknowledging the challenging nature of the information that they are being asked to receive and the decisions that are being made.

In utero transfer to a maternity facility co-located with a NICU should be considered at the earliest opportunity when survival focused management is planned. All such transfers should be discussed with the receiving team, and parents should be made aware that the prognosis (and therefore management) may be revised following *in utero* transfer to a centre with greater experience of managing extremely preterm birth (e.g. following detailed ultrasound scanning). Communication and agreed plans should be documented in full (BAPM Practice Guide: In-utero Transfer) and, when relevant, clearly communicated with the receiving centre. The agreed plan of management should be revised regularly if pregnancy continues. Parents should also be helped to appreciate that the baby may be born in unexpectedly poor, or unexpectedly good condition, and the implications of this for what care might be appropriate. Processes should be in place to ensure timely transfer. Both written and verbal information should be given to parents, with appropriate use of translation services. In some cases, poor maternal health and/or advanced stage of labour may mean that *in utero* transfer is not the safest option.

When survival focused care is planned and time allows, parents should be given an opportunity to visit the neonatal unit and to meet staff and should receive information and support regarding expressing breast milk.

Where appropriate, the practicalities of commencing, withholding and/or withdrawing intensive care and the positive role of comfort focused care strategies should be described to the parents. This will help prepare them for possible outcomes after the birth. Parents may find the access to neonatal psychology services, support and advice from their family, friends, spiritual advisers and/or local and national support organisations to be of great value at this time and should be signposted appropriately.

3. Agreeing and documenting a management plan


Following consultation with parents, initial management of the birth will follow one of two pathways: “Survival focused” or “comfort focused” (Figure 2). Parallel planning can be a useful

strategy to discuss these options. This includes (where desired), providing survival focused care, but discussing the possibility of changing to comfort focused care in the event that a baby responds poorly to attempted stabilisation or develops serious complications.

Consistency in obstetric and neonatal management is essential, both to ensure that the baby is born in the best possible condition and to avoid unnecessary intervention. The agreed plan should be clearly documented and communicated to all members of the obstetric and neonatal teams who may be involved in care of the family.

The challenges inherent in evaluating risk should not be underestimated and categorisation and communication of risk should be undertaken by the most senior clinicians available. Electronic risk calculators may be of value but care should be taken to ensure they are populated with the most recent data and include the most relevant denominator.

Figure 2. Decision-making around management of birth, following risk assessment and after consultation with parents.



Obstetric management

Obstetric management focused on maternal wellbeing and neonatal survival

- When it has been agreed that potentially life-sustaining care for the baby is appropriate, survival focused obstetric care is important to ensure the baby is born in the best possible condition. An individualised package of obstetric intervention should be offered in all cases where a commitment to survival focused neonatal care is in place (David, 2018). This should be in line with the [Saving Babies Lives Care Bundle, Version 3](#) and [BAPM Perinatal Optimisation Pathway](#) and should utilise all optimisation resources, including clinical and parent optimisation passports found in the pathway. The potential for each component intervention to optimise the condition of the individual baby at birth should be considered, and not excluded on the basis of gestational age alone. Obstetric management should be regularly reviewed, particularly if events suggest changing prognosis for the baby. All discussions and decisions should be documented clearly in the clinical record.

The package of obstetric care to be offered to parents *may* (but not necessarily) include any or all of the following:

- **In utero transfer to a tertiary maternity centre with a co located NICU** optimises outcomes for the baby, is better than *ex utero* transfer and is now a prioritised recommendation across all nations of the UK (NHS England Saving Babies Lives Care Bundle (version 3), [BAPM Practice Guide: In-utero Transfer](#)).
- While the majority of women presenting in threatened preterm labour before 27 weeks of gestation do not deliver in the subsequent 24 hours, and transfer may present challenges for the family as well as obstetric and ambulance services, the Working Group strongly recommends that this is considered at the earliest opportunity.
- **Joint obstetric and neonatal counselling** should occur where possible, or at the very minimum, a conversation between the neonatal and obstetric consultant should take place prior to a conversation with the parents. These discussions will inform the decision whether to provide neonatal survival or comfort focused care and obstetric management. If in a non-tertiary centre, this discussion will also inform whether a mother is transferred in utero for the baby to be born in a maternity unit with a co-located NICU.
- A decision for antenatal transfer should be made in a timely manner to facilitate earliest possible in-utero transfer of the mother. Some regions now offer neonatal cot and maternity bed finding services to facilitate in utero transfers. A careful risk assessment should be conducted to ensure that the mother is fit for transfer and that birth in transit is unlikely. Discussions between the referring unit, transport team and receiving tertiary NICU are recommended, and can be facilitated through use of Microsoft® Teams or other conference calling modalities.
- A decision to transfer antenatally does not necessarily commit management to survival focused care and assessment of risk should be an ongoing assessment and reviewed when birth is imminent. Decisions around mode of birth should also be reviewed to ensure appropriate balance of risks and benefits for the mother and baby.
- **Optimally timed course of intrapartum antenatal steroids, tocolysis, magnesium sulphate for neuroprotection.** Optimal timing for cord clamping is uncertain; however deferring clamping the cord for at least one minute (Fogarty, 2018) has been shown to

reduce mortality in preterm infants. Parents should be made aware that there is a paucity of data in relation to the magnitude of benefit and risks of these interventions, particularly below 24 weeks of gestation. (LeMoine, 2025).

- **Maternal Intrapartum Antibiotic Prophylaxis to reduce the risk of neonatal Group B Strep (GBS) disease** – The risk of early onset GBS disease in babies of those women who deliver preterm is estimated to be 2.3 per 1000, with a mortality rate in preterm babies 10 times that of term infants. Current national guidance advises that intrapartum antibiotic prophylaxis should be administered to any woman in established preterm labour (RCOG GTG No 36, NHS England Saving Babies Lives Care Bundle (Version 3)).
- **Intrapartum fetal heart rate monitoring** Below 26 weeks of gestation, a senior obstetrician should be involved in decisions around intra-partum fetal heart rate monitoring as there is a lack of evidence to inform practice (NICE NG25). However, assessment of whether the fetal heart was present before birth by intermittent auscultation or ultrasound is helpful to the attending neonatal team, particularly where survival focused care is planned. The family should be made aware of the rationale for either recommending or withholding fetal heart rate monitoring, this should be discussed and agreed on an individual basis. For example, it may be appropriate not to monitor the fetal heart if birth by caesarean section is not part of the agreed package of care, either because it is considered that the risks of caesarean section outweigh any potential benefits or because parents have declined caesarean section should there be a fetal heart rate abnormality. Autonomic immaturity at gestations below 26 weeks makes interpretation of continuous electronic fetal heart rate monitoring (CEFM) difficult and there is no evidence that CEFM improves outcomes compared to intermittent auscultation. From 26⁺⁰ weeks of gestation, when survival focused management is planned, women in established preterm labour should be recommended CEFM (NICE NG25).
- **Mode of birth** The decision on mode of birth should be informed by the balance of risks and benefits to baby and mother. In the majority of extremely preterm births the mother presents in spontaneous labour and an uncomplicated vaginal birth may be anticipated. The risk of head entrapment following breech presentation is approximately 10% but the evidence for birth by caesarean section for extremely preterm babies is limited and of poor quality (Grabovac, 2018, Reddy, 2015) and prognosis is more likely to be dictated by factors other than mode of birth. NICE guidance is that birth by caesarean section may be considered in cases of breech presentation **after 26 weeks of gestation** (NICE NG25). Extremely preterm caesarean sections can be difficult, and fetal trauma including head entrapment can still occur. Maternal risks and consequences including pain, haemorrhage, infection, thrombosis and injury to bowel and bladder are higher after caesarean section compared to vaginal birth, particularly at extremely preterm gestations, and should be discussed with the mother. There is an impact on future pregnancies in terms of increased risk of uterine rupture and morbidly adherent placenta, and the likelihood of classical caesarean section, with its increased risk of serious maternal complications, is greatest at the most preterm gestations ^(40,41) (Reddy, 2015, Blanc, 2019). For all these reasons it is essential that obstetric care is individualised after full discussion between the family and a senior obstetrician along with the neonatal team. If the baby is deemed extremely high risk for a poor outcome, a decision for a caesarean section should only be made if there are benefits for the mother, or maternal choice following informed counselling of the above risks. There should be clear documentation of this conversation and the mother's wishes (RCPSG Statement). Where birth by caesarean section has been agreed as the optimal mode of birth, this should only occur once labour is established, unless maternal or fetal condition dictates otherwise. Established preterm labour can be difficult to determine and even at

advanced cervical dilation birth may not occur for several days. There may be additional benefit to the baby of delaying birth. In the absence of labour and where birth should be expedited for maternal reasons (e.g. pre-eclampsia or chorioamnionitis) or, more rarely, for fetal reasons (e.g. severe fetal growth restriction) birth by caesarean section may be the only option to ensure timely birth for mother and/or baby. Induction of labour is unlikely to be appropriate in such circumstances where there is maternal or fetal compromise and a commitment to potentially life-sustaining care for the baby has been agreed with the parents.

Obstetric management focused on maternal wellbeing

When a decision is made for comfort focused management of the baby at birth, only interventions for maternal benefit are appropriate. This may include a discussion of options for the pregnancy where there is a risk of maternal morbidity and mortality and high risk of neonatal mortality. This is particularly important in cases of PPROM and chorioamnionitis and should, where appropriate, include input from fetal medicine and preterm birth experts. Intrapartum fetal heart rate monitoring is not advised, although assessing or listening for the presence of a fetal heart to check viability may be helpful in clarifying expectations around the baby's condition at birth and be preferable for parents. Assessment of signs of life after birth should be undertaken following the MBRRACE-UK guidance. (resources including videos are available MBRRACE-UK Guidance, 2020) ensuring that care following birth is respectful and that the individual needs of the baby and parents are prioritised at this difficult time. Parents should be made aware that their baby may show signs of life after birth, including visible heartbeat, gasping and/or movement of limbs but also that some babies who have died before birth may show brief reflex movements.

Neonatal management

Survival focused neonatal care

- This should be in line with the [BAPM Perinatal Optimisation Pathway](#) and should utilise all optimisation resources, including clinical and parent optimisation passports.
- Stabilisation and support for transition should be carried out by, or under the direct supervision of, the most senior member of the neonatal/paediatric team available at the time of birth, and in accordance with **Resuscitation Council UK guidance**, noting specific recommendations for preterm babies (Resuscitation Council UK). Ideally this team will be experienced in stabilisation of extremely preterm babies, including an experienced intubator (BAPM Airway Safety Standard) and led by a consultant neonatologist.
- **Deferred cord clamping for at least 60 seconds** should be routine practice. Perinatal teams should work to develop ways to reliably provide this, in conjunction with thermal care and initial stabilisation, in different birth circumstances. **Normothermia** should be a key area of focus, with the use of a plastic bag and/or other methods of delivering thermal care, and skin protection.
- **Stabilisation and supported transition with lung inflation**, using an appropriately sized facemask, should then be initiated. Care should be taken not to over distend the lungs.
- **Intubation:** Clinical assessment immediately following birth is not a good predictor of survival in extremely preterm babies (Manley, 2010); if there is no response to mask ventilation, and any doubt around the adequacy of ventilation, the baby should be intubated by the most experienced intubator available ([BAPM Airway Safety Standard](#)) and surfactant administered. The most important intervention is establishment of adequate lung recruitment, and the most important measure of success is heart rate.
- Use of **advanced measures for resuscitation** including cardiac massage and endotracheal or intravenous adrenaline are rarely required following extreme preterm birth. In the absence of sufficient evidence to justify a different approach in extremely preterm babies, if advanced resuscitation is considered appropriate, the Working Group recommends applying newborn resuscitation algorithms as used in more mature babies. Clinicians should discuss the possibility of more extensive forms of resuscitation with parents prior to extremely preterm birth and involve them in decisions. In some extremely preterm babies who are already designated to fall into a 'high risk' category, clinicians may consider that the failure to respond to initial measures (including intubation and ventilation) would indicate that the baby now has an 'extremely high risk' of adverse outcome, and that it would be in their best interests to move to comfort focused care (Mactier, 2020).
- Stabilisation should normally be undertaken in the same room as the parents, who should be offered the opportunity to **see, touch and photograph their baby**. Following stabilisation, **contact (for example cuddles)**, may provide positive experiences for parents and can be considered where feasible and appropriate. Following successful stabilisation of the baby, the mother should be supported to express **breast milk** as early as possible, with ongoing facilitation of parental contact and family involvement as partners in care.

Where babies are born in much poorer condition than expected it may be appropriate to reconsider the planned provision of survival focused care and to move to comfort focused care (see [Appendix 2](#) and [3](#)). Absent heart rate or severe bradycardia persisting despite *effective* resuscitation for more than a few minutes is associated with high rates of mortality and neurodevelopmental impairment in extremely preterm babies (Wyckoff, 2012, Haines, 2016). The most senior experienced attending professional should decide if or when attempts to stabilise and/or resuscitate the baby should stop (McGrath, 2016, Wilkinson 2020).

Extreme preterms born out of hospital should have been managed according to the BAPM Framework 'Pre-hospital management of the baby born at extreme preterm gestation'. Deferred cord clamping and mask ventilation via a self-inflating bag is advised and ideally the baby should arrive in a plastic bag, swaddled in a blanket on a thermal mattress. The provision of thermal care and mask ventilation en route does not mean that the hospital team are obliged to provide ongoing survival focussed care following admission to hospital. Management subsequently should be guided by the same principles as described above. For more detail see [Appendix 3](#).

Comfort focused neonatal care

Where there is an extremely high risk of a poor outcome for the baby, it would be in the best interests of the baby to provide comfort focused care, and standard practice not to offer survival focused neonatal management.

The aim of comfort focused neonatal management is to support the parents and their baby and to avoid interventions that may cause discomfort, pain or separation of the baby from the parents. This care should be delivered in the most appropriate location for the family (which is not necessarily a neonatal unit) and should not necessitate *in utero* transfer. There should be an emphasis on family centred care, with opportunities for parents to create positive memories of their baby. (Bliss 2024) An Individualised Care Plan should be made in partnership with parents following guidance (Larcher, 2015) within the BAPM Framework: 'Recognising uncertainty: an integrated framework for palliative care in perinatal medicine'.

Depending on parents' wishes and service provision, a senior neonatologist or paediatrician may be present at birth to provide a brief assessment of the baby's condition at birth and to support midwifery staff and the family. Respiratory support (including provision of positive pressure ventilation) should not be provided. Parents should be offered the opportunity to hold and to spend as much time as they wish with their baby in a quiet and private location. Please refer to the [MBRRACE-UK Guidance 2020 Determination of signs of life following spontaneous birth before 24⁺⁰ weeks of gestation where, following discussion with the parents, active survival-focused care is not appropriate](#). Parents should have been counselled that the baby may show signs of life after birth including visible heartbeat, gasping and/or movement of limbs but also that some babies who have died before birth may show brief reflex movements.

In the rare scenario of the baby being born in much better condition than expected, comfort focused management may need to be reconsidered (see [Appendix 3](#) for further discussion).

Babies born before 24 weeks of gestation who receive comfort care following birth may live for between a few minutes to several hours (Macfarlane, 2003). Supplemental oxygen is not necessary but could be provided if parents desire.

After the baby has died, a parent-led bereavement care plan should be put in place for the family, including communicating with parents and creating memories, referring to the BAPM Framework: 'Recognising uncertainty: an integrated framework for palliative care in perinatal medicine'. Parents should understand what to expect in terms of a review into the care provided during pregnancy and birth using the Perinatal Mortality Review Tool, the national Child Death Review process, and the benefits of investigations such as autopsy and placental histopathology, to provide as much explanation as possible for the preterm birth and the death of their baby (Together for Short Lives). In England, this will include mandatory discussion with local medical examiners (NHS England Medical Examiners System). In England and Scotland this should follow the guidance outlined in the [National Bereavement Care Pathway \(NBC Pathways\)](#). In Wales and Northern Ireland there are locally developed bereavement pathways. Parents should be facilitated to make informed choices

and signposted to support available after they go home. Follow up pathways for all women who have undergone an extremely preterm birth should be in place and include planning care for future pregnancies. Placental histology, undertaken by a perinatal pathologist, should be routine.

After discharge home, optimal communication with all professionals involved (and in particular the GP, health visitor and community midwife) is essential. The mother will continue to require postnatal care and should also receive information and advice about milk suppression or donation (BAPM Framework '[Lactation and loss](#)'). Parents should be offered bereavement counselling and the opportunity to meet with perinatal staff for a follow up consultation in an outpatient setting. Where possible, this meeting should be conducted by the same staff that counselled the family in the peripartum period. Parents should also be offered the opportunity to contribute to a multi-professional perinatal mortality review process that follows the framework set out in published statutory and operational guidance ([NBC Pathways](#) in England and Scotland, 2022). At an appropriate time, the prognosis for future pregnancies should also be discussed.

DRAFT

Implications of this framework for Transport/LNU and SCU teams

Babies less than 27 weeks gestation should be delivered wherever possible in a unit with a NICU. If extremely preterm infants are born in an LNU or SCU, this will increase their risk of poor outcome and potentially affect their risk category in this framework.

For example, most babies born prior to 23 weeks gestation in a maternity unit without a colocated NICU (ie an LNU or SCU) will be **extremely high risk**. UK data indicates extremely low survival rates for babies transferred ex-utero at this gestation. As a consequence, for such babies neonatal management focused on comfort should be provided, and it would usually **not** be appropriate to retrieve and transfer ex-utero.

Implementation of this Framework for Practice

A lead team should be identified in each maternity and neonatal facility with responsibility for implementation, education, and dissemination of this updated Framework for Practice and the accompanying parent information. This is likely to be best placed within the designated Preterm Birth Lead Team (Saving Babies' Lives Care Bundle Version 3).

Links to relevant national documents are provided within this document; these should be highlighted and made easily available within each perinatal facility.

Management of extreme preterm birth and the conversations around this can be exceptionally challenging for staff, so it is recommended that implementation of this Framework into individual units is accompanied by education and training in specific consultation skills. Some guidance is offered in [Appendix 3](#) and a suggested format for parental information is to be found in [Appendix 4](#). Examples scenarios are provided in [Appendix 5](#) for discussion and learning.

Networks need to ensure sufficient resource to cope with the predicted number of extreme preterm deliveries, both actual and threatened, and pathways should be in place to ensure appropriate prioritisation and assessment of women likely to deliver extremely preterm. Following birth, the mother should be accommodated in a maternity facility adjacent to her baby.

Recommendations for audit and research

There is limited research evidence to inform optimal perinatal management of extreme preterm birth, resulting in variation in care and outcomes. In particular, babies born at the lowest gestational ages (<24 weeks) are poorly represented in most research studies, and it remains unclear whether this cohort are embryologically and physiologically different, and require a different approach.

BAPM recommend that neonatal units who care for extremely preterm babies should participate in national audit and research activities, in addition to reviewing their own local outcomes. Specifically, given the small number of babies born at 22 weeks in each unit, a national perinatal approach is necessary to meaningfully build knowledge and experience, to improve care and outcomes ([BAPM Nanopreterm Special Interest Group](#)).

The Working Group propose the following research questions:

- What additional information or tools will help inform decision making at the time of counselling?
- Whether and how are infants born 22-24 weeks different to babies born 25-26 weeks gestation? Should their care approaches be different?
- What are the long-term outcomes and what additional health, educational and social care resources are needed to support families and babies born extremely preterm?
- An international research priority setting partnership identified priorities for extremely preterm babies born less than 25 weeks (Peart, 2025). The most important research question identified was 'what can be done in the neonatal intensive care unit to improve long-term health and developmental outcomes?'. Other important areas for research included antenatal interventions and neonatal care at birth, preventing intraventricular haemorrhages, managing pain, postnatal corticosteroid treatment and supporting families.

Appendix 1: Outcomes for extremely preterm babies: international comparisons

Survival/Mortality

International studies indicate incremental improvements in survival for the most premature babies over the last 1-2 decades. Since the 2019 BAPM framework, an increasing number of multi-centre, population studies and meta-analyses have reported survival figures for babies born extremely preterm including at 22 weeks gestation. Table 1 shows published UK population data (England and Wales) up until 2021 alongside international data; colour coded to indicate whether studies are population based or limited to tertiary and quaternary centres. Even more recent data from 2023 in the UK National Neonatal Audit Programme Report show unchanged survival to 44 weeks post menstrual age for babies at 22 and 23 weeks gestation, with a continued increase in the absolute number of admissions and survivors at 22 weeks gestation until 2023 (NNAP 2023).

International comparisons should be interpreted with caution, given differences in the provision of survival focused care, how survival is reported, the type of neonatal units (e.g. only including inborn babies born in tertiary centres versus all units serving a population), denominator populations, and differences in attitudes and practice. A recent meta-analysis published in 2024 reviewed survival figures for babies born 22-25 weeks gestation. Among high income countries, there was wide variation in survival estimates of live born babies (from 7% (95% CI 5 – 10) at 22 weeks and 49% (43-54)% at 24 weeks of gestation); with higher survival among those admitted to neonatal intensive care (30% (25-36) at 22 weeks, 61% (57-64) at 24 weeks).

Survival in absolute numbers has increased among those born at the lowest gestational ages admitted to neonatal units. In particular, at 22 weeks of gestation, the most recent cohort studies from US, Sweden and Japan indicate that survival to discharge of livebirths range from 24% in US, 39% Sweden, to 46% in Japan. Survival figures are higher among those provided survival focussed care in US (36%) or admitted to neonatal intensive care in Sweden (46%) and Japan (51%). (Table 1). A meta-analysis which focused on babies born at 22 weeks receiving proactive treatment found a pooled prevalence of 29.0% survival (95% CI 17.2-41.6; 31 studies, 2226 babies, 13 countries). Importantly, the studies included births 1991-2016, which reflects a time in some settings when survival focused care may not be commonly practiced, and need to be interpreted with caution (Backes, 2021). The reported survival rates varied greatly among studies and were likely influenced by combining observational data from disparate sources, lack of individual patient level data, and bias in the component studies from which the data were drawn. There may also be differences in population characteristics, as shown by recently published UK data (Smith, 2023), which showed an increase in the proportion of babies born at 22 weeks with less favourable characteristics for survival (e.g. born in the first half of the 22nd week, weighed less than 500g, not born in a tertiary centre, no antenatal steroids).

Table 1* Search criteria

Country	Unit Type	Years of Birth Included	Survival assessment timepoint	Denominator	Survival (95% CI)			
Gestation (completed weeks)					22w	23w	24w	25w
UK (Smith, 2023)	All babies, all units	2020-21	Discharge from neonatal care	Live births	12% (9-16)	40% (36-44)	69% (66-72)	78 % (76-81)
				Survival focused care	21% (15-27)	42% (38-46)	69% (66-72)	78% (76-81)
				Neonatal unit admissions	30% (21-37)	48% (43-52)	72% (69-76)	80% (78-82)
Sweden (Farrooqi, 2023)	All babies, all units	2017-19	1 year of age	Live births	39% (28-50)	67% (57-75)	Not reported	Not reported
				Neonatal unit admissions	46% (34-59)	70% (61-78)	Not reported	Not reported
Japan (Kono, 2018)	Inborn, selected NICUs	2008-12	Discharge from neonatal centre	Live births	46% (40-52)	73% (71-77)	85% (82-87)	Not reported
				NICU admissions	51% (45-58)	75% (72-78)	85% (83-88)	Not reported
USA (Edwards, 2024)	Inborn, selected NICUs	2020-22	Discharge from hospital	Live births	24% (23-26)	52% (51-54)	70% (69-71)	81% (80-82)
				Postnatal life support	36% (34-38)	55% (54-56)	72% (71-73)	82% (80-82)
Meta-analysis (Li, 2024)	High income countries	2000-2020	Pooled survival (discharge or at 1-3 years old which ever was latest follow up assessment)	Live births	7% (5-10)	26% (22-31)	49% (43-54)	68% (63-72)
				NICU admissions	30% (25-36)	44% (41-48)	61% (57-64)	74% (70-77)

Key

- National level data
- Inborn tertiary/quarternary level NICUs only
- Meta-analysis data

* International Survival Data Search Strategy

A search using Pubmed was conducted including English language results between 2018 and 2025 using the search terms below (in Title/Abstract). Each country/ network was then added to the search to look for national data. No limitation to the type of study was applied.

22 OR 23 OR 24 OR 25 OR 26 AND week(s)

OR extreme(ly) preterm OR premature* OR periviable OR limit viability

AND survival OR death OR mortality OR outcome(s)

Then AND 'Country' or 'Vermont Oxford Network' or VON or 'National Institute of Child Health and Development' or 'NICHD' or "systematic review"

Confidence intervals where not given by papers were calculated using figures using online CI calculator. Kohn MA, Senyak J. Sample Size Calculators [website]. UCSF CTSI. 28 September 2024. Available at <https://www.sample-size.net/> [Accessed 01 October 2024]

UK figures survival for babies born extremely preterm

The latest survival and outcomes data from births in England and Wales for years 2020-2021 are presented in Table 1. Survival has increased steadily and survival focused care has also increased and is provided to 94% (CI 93-96) of babies at 23 weeks and 42% (CI 37-47) of live births at 22 weeks of gestation. (Smith, 2023). Survival at 22 weeks of gestation is based on small numbers of babies and thus the confidence limits are wider than at other gestational weeks.

DRAFT

Table 2. England and Wales births and outcomes born 22-26 weeks gestation

	Births in 2020-2021				
Gestational Week	22 weeks	23 weeks	24 weeks	25 weeks	26 weeks
All births	727	892	993	1081	1403
Births alive at onset of labour	477	655	751	833	1169
Live births	319	559	700	802	1144
As a % of births alive at onset of labour	66.9% (64.1 to 69.8)	85.3% (83 to 87.6)	93% (91.3 to 94.7)	96.3% (95.1 to 97.5)	97.9% (97.1 to 98.7)
Receiving active care/ survival focused care	183	528	700	800 Incl 3 intrapartum still births	1139 Incl 1 intrapartum stillbirths
As a % births alive at onset of labour (95% CI)	38.4% (34 to 42.7)	80.6% (77.6 to 83.6)	93.2% (91.4 to 95)	96% (94.7 to 97.3)	97.4% (96.5 to 98.3)
As a % of live births	57.4 (51.9 to 62.8)	94.4 (92.6 to 96.4)	100.0 (100.0 to 100.0)	99.4 (98.9 to 99.9)	99.5 (99.1 to 99.9)
Admitted for neonatal care	134	467	669	785	1121
As a % births alive at onset of labour	28.1% (24.1 to 32.1)	71.3% (67.8 to 74.8)	89.1% (86.9 to 91.3)	94.2% (92.7 to 95.7)	95.9% (94.8 to 97)
As a % of live births	42% (36.6 to 47.4)	83.5% (80.5 to 86.6)	95.6% (94 to 97.1)	97.9% (96.9 to 98.9)	98% (97.2 to 98.8)
As a % of babies receiving survival focused care	73.2% (66.8 to 79.6)	88.4% (85.7 to 91.2)	95.6% (94 to 97.1)	98.1% (97.2 to 99)	98.4% (97.7 to 99.1)
Survival to discharge	39	222	483	626	966
As a % births alive at onset of labour)	8.2% (5.7 to 10.6)	33.9% (30.3 to 37.5)	64.3% (60.9 to 67.7)	75.2% (73 to 77.4)	82.6% (80.8 to 84.4)
As a % of live births	12.2% (8.6 to 15.8)	39.7% (35.7 to 43.8)	69.0% (65.6 to 72.4)	78.1% (75.9 to 80.4)	84.4% (82.6 to 86.2)
As a % of babies receiving survival focused care	21.3% (15.4 to 27.2)	42.0% (37.8 to 46.3)	69.0% (65.6 to 72.4)	78.3% (76.1 to 80.6)	84.8% (83.1 to 86.6)
As a % of those admitted to neonatal units	33.3% (18.5 to 48.1)	47.5% (43.0 to 52.1)	72.2% (68.8 to 75.6)	79.7% (77.5 to 82)	86.2% (84.5 to 88)

Source:NNRD and MBRRACE-UK data

Adapted table (Smith, 2023) with additional data on births at 25,26 weeks from NNRD and MBRRACE-UK.

Severe impairment

Individual perception of the impact of impairment on functioning in society (i.e. disability) is highly personal and varies from family to family, dependent on their experience, knowledge and attitudes, and the support available to them. Indeed, it also varies between neonatal health professionals (Gallagher, 2001, Gallagher, 2016). What for one individual or family may be an acceptable outcome may not be so for another.

The severe impairment category defined by 2008 BAPM Working Group includes any of:

- Severe cognitive impairment with an IQ lower than 55 (< -3 standard deviation); this will usually result in the need for special educational support and require supervision in daily activities.
- Severe cerebral palsy – classified as Gross Motor Function Classification System (GMFCS) grade 3 or greater. GMFCS grade 3 is walks with assistive mobility devices indoors and outdoors on level surfaces, May be able to climb stairs using a railing, may propel a manual wheelchair; may require assistance for long distances or uneven surfaces.
- Blindness or profound hearing impairment.

Two-year neurodevelopmental outcome data for recent cohorts

The previous BAPM 2019 framework presented prevalence of severe neurodevelopmental impairment assessed using standardised Bayleys assessment for the EPICure2 cohort born in 2006. However, more recent national data show that only 40% of babies born less than 30 weeks in England and Wales have a standardised e.g. Bayleys or Griffiths neurodevelopmental assessment conducted (van Blankenstein, 2024). National data show that of 41 505 babies born less than 30 weeks gestation over the 11-year period 2008-2018 in England and Wales, 24 125 (58%) had a 2-year neurodevelopmental assessment recorded. Completeness improved from 32% in 2008 to 71% in 2018; completeness was around 70% 2014-2018. The majority ($\geq 87\%$) of babies were assessed using functional versus standardised neurodevelopmental assessments. Whilst there are recognised limitations to non-standardised neurodevelopmental assessments, they provide meaningful functional outcomes of babies receiving contemporary neonatal care.

Functional assessments comprised parent reported outcome measures and health professional assessment in the following neurodevelopmental domains: visual, auditory, communication and neuromotor, each graded as 'impairment' or 'no impairment'. In addition health professionals provide an overall developmental outcome; 'normal' (< 3 months delay), 'mild' (3–6 months delay), 'moderate' (6–12 months delay) and 'severe' (> 12 months delay).

Table 3 reports the incidence of 'severe impairment' based on the definition "unable to walk without assistance or blind or hearing impairment not correctable" following a re-analysis of data from a published study that maps to this definition (van Blankenstein, 2024). The risk of severe impairment increases with increasingly preterm birth and is around one in seven at 24 weeks of gestation, and one in four at 23 weeks of gestation for those babies born who receive survival focused care and survive. Rates of impairment were higher for the earlier gestation group across all domains, and there was no improvement over time. Relatively few babies born at 22 weeks of gestation have available long term outcome data, which means estimates of rates of outcomes are imprecise. However, evidence from other studies show that the proportion of babies born at 22 weeks with severe impairment is at least 30%.

UK data informed revised estimates of survival outcomes (Table 2 2008-2018 birth cohort)

22⁺⁰ - 22⁺⁶ weeks: 1-in-3 survivors has severe impairment
23⁺⁰ - 23⁺⁶ weeks: 1-in-4 survivors has severe impairment
24⁺⁰ - 24⁺⁶ weeks: 1-in-5 survivors has severe impairment
25⁺⁰ - 25⁺⁶ weeks: 1-in-9 survivors has severe impairment.
26⁺⁰ - 26⁺⁶ weeks: 1-in-12 survivors has severe impairment.

Table 3
(Adapted from van Blankenstein, 2024)

Gestation (complete d weeks)	Severe developmental delay (over 12 months) n % (95% CI)		Unable to walk without assistance n % (95% CI)		less than 5 words n % (95% CI)		Blind or only sees light n % (95% CI)		Hearing impairment not correctable n % (95% CI)		Severe impairment based on domains (any of vision blind, uncorrected auditory impairment, unable to walk without assistance)	
	N with data	Impairment	N with data	Impairment	N with data	Impairment	N with data	Impairment	N with data	Impairment	N with data*	Impairment
23	527	120 22.8% (19.4 to 26.5)	576	104 18.1% (15.1 to 21.4)	554	175 31.6% (27.9 to 35.6)	552	7 1.27% (0.62 to 2.59)	549	16 2.91% (1.80 to 4.68)	547	114 20.8% (17.7 to 24.4)
24	1404	198 14% (12 to 16)	1542	216 14% (12 to 16)	1492	366 25% (22 to 27)	1471	11 0.7% (0.4 to 1.3)	1452	26 1.8% (1.2 to 2.6)	1438	234 16.3% (14.5 to 18.3)
25	2007	186 9.3% (8.1 to 11)	2241	240 11% (9.9 to 12)	2189	408 19% (17 to 20)	2140	14 0.7% (0.4 to 1.1)	2132	21 1.0% (0.6 to 1.5)	2101	263 12.5% (11.2 to 14.0)
26	2873	215 7.5% (6.6 to 8.5)	3255	267 8.2% (7.3 to 9.2)	3177	456 14% (13 to 16)	3146	16 0.5% (0.3 to 0.8)	3125	33 1.1% (0.7 to 1.5)	3078	292 9.49% (8.5 to 10.6)

[CI calculated using Wilson]

*Only 20 22w children followed up at 2 years hence not included in the table

**Adapted table from van Blankenstein, 2024, CI calculated by the study authors

International neurodevelopmental outcomes in preterm born babies

A search using Pubmed was conducted including English language results between 2018 and 2025. Each country was then added to search for national data. No limitation to the type of study was applied. We did not include single centre or regional studies. Definitions of 'severe NDI' can vary between assessments and reports so care should be taken when interpreting these figures, especially where there are small numbers of patients assessed.

Severe neurodevelopmental impairment at 22 weeks ranged from 36% to 52% among babies born at 22 weeks, and 21% to 42% for babies born at 23 weeks.

Table 4

Country	Unit types	Method of assessment	Years	Assessment timepoint	Level of NDI Reported	Neurodevelopmental Impairment (NDI) in Assessed Survivors (95% Confidence Interval)				
						22w	23w	24w	25w	26w
Sweden [§] (Serenius 2013)	All babies, all units	BSID III	2004 - 2007	30 months	Severe	N = 5 40% (5-85)	N = 47 21% (11-36)	N = 86 13% (7-22)	N = 151 10% (5-16)	N = 167 7% (4-12)
Japan [¶] (Kono 2018)	Inborn, selected NICUs	GMFCS and KSPD	2008 - 2012	36-42 months	NDI*	N = 77 52% (40-63)	N = 308 42% (36-48)	N = 447 34% (30-39)	-	-
US* (Bell 2022)	Selected academic NICUs (NICHD)	BSID III	2013 - 2016	22-26 months	Severe	N = 29 31% (15-51)	N = 278 34% (28-40)	N = 539 29% (25-33)	N = 695 18% (15-21)	N = 878 15% (12-17)
Meta-analysis (Myrhaug, 2019)	High income countries	BSIDII or III	1994 - 2017	18-36 months	Severe	N = 44 36% (24-51)	N = 316 22% (12-38)	N = 968 19% (11-31)	N = 144 1% (10-19)	N = 168 3% (4-18)

Footnotes on NDI Categorisation

[§]Reporting severe NDI, defined as any of BSID III composite cognitive, language or motor score <55 (<3SD), severe cerebral palsy unable to walk even with an aid, bilateral blindness or deafness.

[¶]Grouped neurodevelopmental impairment (not 'severe' NDI). NDI defined as cerebral palsy with ability to walk using aids or more severe (GMFCS >2), visual impairment, hearing impairment, DQ <70 on KSPD (= BSIDIII <85/ <1SD) or delay assessed by physicians where child not fully evaluated.

*Severe NDI, defined as BSID III cognitive composite or motor composite score <70 (<2SD), limited walking even with an aid (GMFCS >3), bilateral blindness or bilateral severe functional hearing impairment.

N is number of children evaluated (not including those lost to follow up). NDI Neurodevelopmental Impairment, BSID Bayley Scales of Infant Development, GMFCS Gross Motor Function Classification System, KSPD Kyoto Scale of Psychological Development, DQ Developmental Quotient. CIs calculated by framework authors where not provided in individual publication(s).

CIs were calculated using Online CI calculator. Kohn MA, Senyak J.

Key



National level data



Inborn tertiary/quarternary level NICUs only



Meta-analysis data

Search terms: Extreme(ly) preterm OR extreme prematurity OR extremely premature OR limit viability OR periviable
OR 22 week* OR 23 week* OR 24 week* or 25 week* or 26 week*
AND neurodevelopment* OR developmental disability* OR mortality OR survival OR child development OR cerebral palsy
OR Bayley scale OR gross motor function classification OR gmfcs OR cognition Or hearing loss Or vision disorders

DRAFT

Appendix 2: Example case scenarios

C1. Counselling at 22+6 weeks, unfavourable risk factors, maternity unit with co-located NICU

A mother presents to a maternity unit co-located with a NICU at 22+6 weeks gestation in imminent labour. She had ruptured her membranes three weeks earlier at 19 weeks gestation, with subsequently oligo/anhydramnios. There is maternal pyrexia, with suspected chorioamnionitis and there has been insufficient time to give antenatal steroids before the baby is born. She has a singleton male baby with an estimated weight of 490g.

Recommended considerations and actions: Whilst this baby is anticipated to be born towards the end of week 22, added unfavourable risk factors place this baby at extremely high risk of death or severe impairment if born today. Comfort care should be provided.

Outcome: The infant is born later that day (birth weight 530g). Paediatricians are not present at the birth. Only brief reflex movements are observed in the first minute, and consequently this is not registered as a live birth (ref Signs of Life). His mother is supported by bereavement midwives on the postnatal ward.

C2. Counselling at 23+6 weeks, favourable risk factors, maternity unit with co-located NICU

A woman is transferred to a maternity centre with a co-located NICU at 23⁺⁶ weeks of gestation in preterm labour following premature rupture of membranes two days earlier. She had received steroids prior to transfer. The female baby appears well grown and there are no signs of fetal compromise. After discussion with the neonatal team, and being informed about the outcomes of preterm birth, the mother expresses that she is very concerned about the possibility of the baby surviving with severe disability. She requests no survival focused obstetric management, and comfort focused care of the baby at birth.

Recommended considerations and actions: The neonatal team should explore the mother's concerns and provide information about the nature and risks of severe disability for her baby. They should advise the mother that, taking all factors into account, her baby would have a moderate risk of dying or of severe disability. It would be usual to provide survival focused care of the baby in this situation, with the knowledge that if complications develop in the neonatal intensive care unit, there would be the option of later withdrawal of life-prolonging treatment.

Outcome: The mother agrees to this plan and the baby is born a few hours later, receives stabilisation in the birth room and is transferred to the neonatal intensive care unit.

C3. Counselling at 24 weeks, fetal growth restriction, local neonatal unit

A mother has been admitted to a local maternity unit (LNU) in preterm labour at 24⁺³ weeks of gestation. The singleton male baby is very small with an estimated weight of 450 grams. The mother has not yet received antenatal steroids. The on-call paediatric team is asked to provide counselling and attend the birth.

Recommended considerations and actions: As per the Framework, the first step is to assess the risk for the baby if birth occurs. At a gestation of 24⁺³ weeks, the average survival rate for liveborn babies in the UK (if survival focused treatment is provided) would be approximately 60%, with a 1 in 7 risk of severe impairment among survivors. However, in this case, the very low birth weight for the

gestational age in a male baby increases the risk. It is difficult to quantify this risk, but the baby's prognosis is worse than average for 24 weeks of gestation, and within the "high" risk category. Given the risk for the baby, counselling should, if possible, be provided by an experienced senior trainee or consultant neonatologist in conjunction with the obstetric team. If possible, this counselling should take place after the local team has discussed the case with the nearest NICU. It would be appropriate to provide survival focused obstetric and neonatal management if that were desired by the parents. However, it would also be appropriate to provide comfort focused care, if that was felt by the parents to be in the baby's best interests.

Outcome: In this case, the parents decide after consultation that they wish the baby to receive comfort focused care. Labour progresses and a baby is delivered showing signs of life and weighing 460 grams. He is bruised and floppy with a heart rate of 50 beats per minute. The paediatric team attend to support provision of comfort focused care. The baby is wrapped and given to his parents to hold. He dies at approximately 30 minutes of age.

C4. Counselling at 22+0 weeks, SCBU and comfort care

A mother presents to her local maternity unit (SCBU) at 22⁺⁰ weeks of gestation with bulging membranes and active preterm labour. No antenatal steroids have been given, and the estimated weight of the male baby is 510g.

Recommended considerations and actions: In this situation, there is an extremely high risk of poor outcome for this baby if birth occurs within a short period of time. The mother and her partner should be informed of the likely outcome, and advised that if baby delivers in the near future (as appears likely) intensive medical treatments would not help baby. If baby does deliver in the next day or two, baby will be provided with comfort focused care.

The baby's father asks if there isn't anything the neonatal team could do to save baby.

The neonatal team should sensitively explain that if baby arrives today it is too early to save baby. However, if birth does not occur imminently, and the baby stays in utero for even a few days, it may be possible to consider antenatal steroids and transfer to a hospital with a NICU. In that situation, the risk for the baby would still be very high, but survival focused neonatal care may be possible.

Outcome: Labour progress in the next 24 hours. The mother receives obstetric management focused on maternal wellbeing, and the baby is born with no signs of life.

C5. Counselling at 22+3 weeks, labour without chorioamnionitis, in utero transfer from LNU to NICU

A mother presents in preterm labour to a maternity centre without a co-located NICU (eg a Local Neonatal Unit or Special Care Unit setting) at 22⁺³ weeks of gestation. The baby was conceived by IVF and gestation is certain. The baby is female and has an estimated fetal weight of 480 grams. The mother has not yet received antenatal steroids. The paediatric team is asked to provide counselling and attend the birth.

Recommended considerations and actions: Risk assessment in this case indicates that if birth occurs imminently, there would be an extremely high risk of the baby dying or of surviving with severe impairment. If labour progresses, it would be usual to provide comfort focused care at birth. The neonatal team may attend the birth to provide support for comfort focused care, but not to provide resuscitation. However, there are potentially modifiable risk factors in this case. In the absence of evidence of chorioamnionitis, it may be possible to delay preterm birth with tocolysis and so, if the

parents desire, an survival focused approach to management, antenatal transfer and corticosteroids could be provided.

Two days later, at 22⁺⁵ weeks, following transfer to a maternity unit co-located with NICU and administration of steroids, labour progresses.

Given the advance in gestation, availability of specialised neonatal intensive care, and anticipated effect of corticosteroids, the baby's risk is now judged to fall in the "high" category. Accordingly, after further consultation with parents, it would be appropriate to provide survival focused management if this is what parents wish.

Outcome: In this case, the parents decide that they wish the baby to receive survival focused neonatal care, magnesium sulphate is given, and a baby is born showing signs of life and weighing 490 grams. The baby is intubated, receives surfactant and is transferred to neonatal intensive care.

C6. Counselling 25+2 and declines in utero transfer to NICU

A mother presents to her local maternity unit (SCBU) at 25⁺² weeks of gestation in early labour. She has a well grown female baby. She is offered, and accepts, antenatal steroids, but declines transfer to the nearest NICU (1 hour's journey away) where a cot is available. Her partner asks the obstetric team what would be best for the baby, and it is clear to the midwife that parents disagree about transfer.

Recommended considerations and actions: As part of the consultation with parents, the reasons why transfer is being recommended should be clearly explained. Data show that for the most preterm babies, prognosis (both survival and neurodevelopmental outcome) is better if they are delivered in a maternity unit adjacent to a NICU. It can be difficult to predict preterm labour, and so early transfer is preferred. It would be important to explore reasons why the mother does not wish to be transferred and to address all of her concerns. If the mother still refuses transfer she cannot be moved, but she should be offered magnesium sulphate and counselled that the baby will be moved after birth. In this instance, the best interests of the child would be served by early care in a NICU. It would be prudent to alert both the NICU and the local transport team, as well as to think about who will be available to stabilise the baby after birth. The mother should also be informed that, while every effort will be made to move her to a maternity facility adjacent to the NICU, her transfer may need to be delayed if she is unwell after birth.

Outcome: The baby is born 24 hours later in the SCBU. The local team provide survival focused care, including thermal management and non-invasive respiratory support. The local transport team are called, and transport the baby to the closest NICU.

Appendix 3: Situations of uncertainty and potential conflict: recommended considerations and actions

S1. Uncertain gestational age

If gestational age is uncertain, (*i.e.* no dating ultrasound scan) but thought to be $\geq 22^{+0}$ weeks, an ultrasound scan by an experienced sonographer should be carried out if time permits. If the fetal heart is heard during labour, a professional experienced in stabilisation of extremely preterm babies should attend the birth. The baby should be delivered into a plastic bag and an estimate made of gestation. Unless the baby is clearly $< 22^{+0}$ weeks of gestation, and/or estimated (or weighed) at < 350 g, stabilisation and supported transition with lung inflation, using an appropriately sized facemask, should begin, usually after one minute of deferred cord management. Subsequent management will be dictated by the clinical condition of the baby, the response to stabilisation manoeuvres and parental wishes and expectations. In this scenario, it is likely that the parents will have had little, if any, time to consider the situation and so it may be appropriate to proceed with initiating survival focused neonatal care and to reassess the situation in the ensuing minutes, hours and days. It is noted that using condition at birth to assess either gestation or risk of poor outcome is not reliable (Manley, 2010).

S2. Rapid birth without time for counselling

Preterm labour often progresses rapidly, and there may be insufficient time for detailed discussion with the parents before the baby is born. In such a scenario, a decision about management at birth will need to be made based on the available clinical information and informed by the most recent management plan, if any. When risk is unclear (for example gestation is uncertain), and particularly if there has not been time for full discussion with parents, it would usually be reasonable to embark on a provisional plan of stabilisation +/- resuscitation – providing potentially life-sustaining treatment at birth, but redirecting to comfort focused care if the baby appears very immature or responds poorly to stabilisation (for example remains severely bradycardic despite intubation and intermittent positive pressure ventilation). Some extremely preterm babies will present to hospital after having been born in the out-of-hospital setting and having been managed according to the relevant [BAPM pre-hospital framework](#). (See also [Appendix 2 C6](#))

S3 Baby born in unexpectedly good condition

In the rare circumstance where comfort focused care has been agreed, but a baby is born in unexpectedly good condition, attending midwifery and/paediatric medical staff should discuss with parents whether the estimated gestation and prognosis were accurate and whether the planned comfort focused approach is still appropriate. Stabilisation should not be delayed if deemed in the baby's best interests.

S4. Baby born in unexpectedly poor condition

When survival focused neonatal care has been agreed, but the baby is born in unexpectedly poor condition, it is the responsibility of the most senior attending neonatal professional to decide if ongoing attempts at stabilisation and/or resuscitation are in the baby's best interests. This should be conveyed sympathetically but unambiguously to parents, and comfort focused care offered.

S5. Parents requesting survival focused care in settings of extremely high risk.

Where preterm birth has been assessed to be at extremely high risk of poor outcome, comfort focused care would be appropriate and survival focused care should not be offered. For example, this would apply to an baby being born at the start of the 22nd week of gestation, in a centre without a co-located NICU and without antenatal steroids.

In such situations, if parents request or insist on survival focused care, attending neonatologists should sensitively explain that sadly it appears that baby is arriving too early. If birth is able to be delayed for even a few days, other options may be available. (see [Appendix 2](#), c2, c3)

If the parents wish, they should have the opportunity to discuss outcomes with a second senior member of the perinatal team. When parents do not agree with the perinatal team, recently published RCPCH guidance around dealing with conflict may be helpful (RCPCH 2019).

S6. Threatened birth before 22⁺⁰ weeks of gestation

Where gestational age is certain and is below 22⁺⁰ weeks, it would not be considered in the best interests of the baby, to offer survival focused care. If birth is imminent, there would be no benefit (to the baby) in arranging transfer of the mother to a centre with a co-located NICU. If it is possible that the birth may be delayed to a point where survival focused care of the baby would be planned, transfer of the mother to a maternity unit adjacent to a neonatal intensive care unit should be considered.

S7. Born outside of hospital

If a maternity unit receives a call regarding a woman in preterm labour who is imminently delivering or if a preterm baby has been born prior to arrival at hospital the caller should be recommended to call an ambulance.

The ambulance clinicians will follow JRCALC (Joint Royal Colleges Ambulance Liaison Committee) guidelines to provide care to the woman and baby. These guidelines are aligned to the BAPM Framework: “Pre-hospital management of the baby born at extreme preterm gestation”. Where gestation is uncertain, ambulance clinicians should begin to provide survival focused care (see [S1](#)). The prehospital guidelines closely reflect the content of this document, but interventions will likely be limited to facemask ventilation with a self-inflating bag and thermal care with a plastic bag, thermal mattress, swaddling and foil blanket. A pre-alert should be placed to the receiving hospital, as directed by their local policies.

Hospital staff receiving babies from the ambulance service should be prepared to greet the ambulance crew immediately and assess the baby on arrival. Further management should be in accordance with the principles in this framework based on a full assessment of the situation and the baby's condition on handover. The provision of thermal care and mask ventilation en route does not mean that the hospital team is obliged to provide ongoing survival focussed care following admission to hospital. Birth outside hospital will often be associated with significantly increased risk of mortality and morbidity. Where it is clear following assessment that the baby is at extremely high risk of poor outcome, this should be sensitively communicated to parents and the baby should be provided with comfort focused care. Should the mother be travelling separately to her baby, care should be continued pending her arrival.

On occasion, a prehospital critical care team may be involved, and in these cases interventions may include intubation. Networks should liaise with teams providing this care to establish pathways for providing remote support and guidance.

It should be recognised by hospital staff that for ambulance personnel this will be an extremely rare occurrence and a situation with which they are unfamiliar. As such it will be a profoundly stressful situation for them and care should be taken to ensure that communication is respectful and sensitive. Ambulance clinicians and emergency department staff should not be expected to counsel parents regarding likely outcomes and options regarding ongoing management; this would be the role of the neonatal, obstetric and midwifery teams. Pre hospital clinicians should be encouraged

and supported to attend the Out of Hospital NLS (OHNLS) courses available through the Resuscitation Council UK to gain further expertise in this area.

S8. Live birth following medical termination of pregnancy

Rarely, neonatal teams may be called by the maternity teams to attend an extremely preterm baby (for example at 22 or 23 weeks gestation) who is liveborn following a medical termination of pregnancy. It is important to clarify the reason for requesting attendance of the neonatal team. It is preferable for the baby to be seen by a member of the obstetric team, rather than the neonatal team to avoid confusion and to support notification to the coroner (required for all live births following termination of pregnancy) (RCOG Position Statement 2023). A [National Bereavement Care Pathway for Termination of Pregnancy for Fetal Anomaly](#) (also referred to as 'medical reasons') is also available for reference (NCBP ToPFA, 2022).

If a neonatal team does attend the birth, the same principles for care of the baby apply as to any extremely preterm birth. Because of the gestation of such births (almost all will fall into the Extremely High Risk category) and the nature of parental wishes, comfort-focused care should be provided.

DRAFT

Appendix 4: Communication: Guidance for professionals consulting with families at risk of extreme preterm birth.

This Appendix is designed for use by all staff caring for families at risk of extreme preterm birth, to facilitate the sharing of consistent and accurate information. It should be used to support conversations about decision making with parents, in conjunction with written information such as the suggested *Template Parental Information - Helping parents to understand extreme preterm birth* (Appendix 4). Written information should *never* be used as a stand-alone information sharing tool. In order properly to involve parents as equal partners in care and decision-making for their babies, all parents facing potential extreme preterm birth need to understand the risks associated with their baby's birth, and possible treatment options.

In the context of the different risk scenarios outlined in this Framework, health professionals need to consider how their approach to consulting with parents may differ, depending on the individual circumstances:

- In **extremely high risk** cases, parents should be provided with relevant information about the risk to their baby, and the recommendation that it would be best for their baby to provide obstetric care focused on maternal wellbeing and comfort focused neonatal care. Parents should be told about the role they can play in caring for their baby and memory making after birth, and should be fully involved in decisions about how and where comfort focused care takes place.
- In **high risk** cases, the role of the consultation is critical in supporting parents together with professionals to decide on the right pathway for their baby. Parents should be provided with as much information as is available and should have as much time as possible, and where possible, over a number of discussions, to work through the different options available in order to agree with professionals what the right option is for their family.
- In **moderate risk** cases, parents should be provided with relevant information about the risk to their baby, and the recommendation that it will be best for their baby to provide survival focused care both antenatally and after birth. Parents should be told about likely interventions and what may happen next, and be fully involved in decisions about how survival focused treatment is managed.

Thus, consultation is most critical where birth would be associated with a high risk of a poor outcome (see main text). In such situations, parents need support to make an informed choice about the provision of either survival focused or comfort focused management; such situations demand the greatest care and sensitivity. Consultation should not be directive, but professionals should seek to determine when gentle guidance around what is likely to be in the baby's best interests would be helpful for the family.

When is the right time?

Evidence suggests that parents find that, where time allows, consultation is most useful at the earliest opportunity, both to allow time for information processing, discussion, and decision making, and to minimise the effects of labour and medications on cognition. Follow up consultations allowing ongoing dialogue are highly valued by families (Young, 2012, Kharrat, 2018) and should be offered at any point, acknowledging the challenging nature of the information that parents are being asked to receive, the time this may take to process and the decisions that need to be made.

Who should be involved?

Consultation with parents should ideally be provided by the most experienced members of the perinatal team involved in care of the mother and her baby. Continuity of care is essential and, whenever possible, consultation should be delivered as a joint obstetric, neonatal and midwifery approach, ensuring transparency and consistent, clear communication. The presence of members of the multidisciplinary team (particularly nurses and midwives) during such conversations is highly valued by families, and may provide opportunity for clarification and ongoing conversation outside the formality of such settings (Kharrrat, 2018, Kaempff, 2009). Parents may also find the advice from psychology professionals, and support of their family, friends, spiritual advisers and/or voluntary organisations to be of great value at this time.

Structuring the consultation

1. **Exploring the parents' prior knowledge and understanding** can be a useful way to open the consultation. Establishing parents' own understanding about the risks of their situation, their prior experience and knowledge, as well as their expectations of the conversation is important, both to generate trust and to ensure that the consultation meets their individual needs. Parents' hopes, priorities and expectations of the care that they and their baby will receive should be explored with sensitivity, honesty and compassion in a realistic way.
2. **Balanced Information** - Studies suggest that conveying solely negative information to parents is not well received. Providing balanced information with honesty seems to be most useful to parents. Respecting parents' perspectives and the importance of hope, even in the most difficult of situations, is highly valued. Exploring parental hopes, wishes and fears in each scenario can help to do this, and to build trust and rapport with the clinical team. Where survival is not possible, or is extremely unlikely, parental hopes relating to spending time with their baby, involving family members, and memory making should be explored.
3. **Conveying Risk** - Categorisation of risk to the baby of death or survival (with or without impairment) in a given scenario should be conveyed sympathetically and with clarity. Parents may find it useful to see this displayed graphically (see [Appendix 5](#)). Gestation-based risk should be explained within the context of other risk modifiers (such as birth weight, gender, multiplicity, etc.). It is important to convey information accurately, in the appropriate context. While the most relevant statistic for parents is usually the chance of survival if survival focused stabilisation and neonatal intensive care is attempted, parents should be helped to understand that not all babies survive labour, and so outcome data depend upon the stage at which parents are being counselled. Not all parents find percentage figures easy to understand. It can be helpful to explain in terms of odds e.g. 1 in 4, or 1 in 10. To avoid framing bias, we suggest interpreting risk neutrally. For example, *"Given what we know about the situation for your baby, there is a 30% chance of your baby surviving. This means that for every 10 babies treated with intensive care in situations like this, three would survive while sadly seven would not"*.
4. **Discussing Poor Outcomes** - There is not a simple definition of a 'poor' outcome – the interpretation of this is likely to vary greatly between clinicians, parents, and families. Published data generally refer to scoring systems and classification of motor and cognitive dysfunction, but also often include children with profound vision or hearing loss. Some of these terms may not be meaningful to families, and families' views may differ on the outcome that they would regard as unacceptably poor. Therefore, discussions should always include exploration of the parents' views and values relating to an acceptable outcome. Conveying the concept of severe disability in childhood, and the possible implications for future quality of life, can be difficult. Guidance can be found in [Appendix 5](#). And refer to BAPM framework of Palliative care Uncertain Future
5. **Discussing Comfort focused Care** – Where appropriate, the practicalities of commencing, withholding and withdrawing intensive care and the positive role of comfort focused care

should be described to the parents. This will help prepare them for possible outcomes after the birth. Signs of life may or may not be present after birth. If present, the length of time that the baby may show signs of life will vary (MBRRACE-UK). It can be useful to speak about memory making, exploring parents' hopes and wishes. We suggest referring to the BAPM Framework: 'Recognising uncertainty: an integrated framework for palliative care in perinatal medicine'.

6. **Decision making** – A shared decision making process is vital, especially in situations of moderate to high risk of unacceptably poor outcome. Support and guidance should be tailored to the needs of each family. Parents should be helped to understand that, even taking all available information into account, babies may be born in unexpectedly poor or unexpectedly good condition, and that this may impact upon what care at birth would be best for their baby.
7. **Parental involvement in care** - Evidence suggests parents find it very useful to hear how they can be involved in care for their baby. Neonatal care in the UK now typically implements Family Integrated Care (FICare), a model and philosophy of care within which families are enabled to be primary caregivers to their babies in partnership with clinical teams. This is proven to be of benefit to babies and parents, including in cases of extremely premature birth. Units can support families in this model of care by 'listening to them, building on their strengths, and encouraging their participation in experiences and decision-making to enhance control and independence.' ([BAPM FICare Framework for Practice](#); [Bliss Baby Charter](#)) or to view a video tour of the unit if one is available.
8. Where it is planned to offer survival focused care to the baby, and time allows, parents should be given an opportunity to visit the neonatal unit and to meet staff or view a video tour of the unit if one is available. Parents should receive information and support regarding expressing breast milk and the other ways that they could be involved in the hands-on care of their baby if s/he is admitted to the neonatal unit.
9. **Documentation and follow up** - Communication and agreed plans should be documented in full in the clinical record and plans revised regularly if pregnancy continues and/or depending upon the condition of the baby at birth and in the early days after birth. If *in utero* transfer is undertaken, the content and results of previous conversations should be clearly communicated (verbally, and in writing) with the receiving centre. Evidence suggests that parents find it very useful to receive supplemental information such as written information, visual aids and links to other resources. We include a suggested template for this information below.

Appendix 5: Helping parents to understand extreme preterm birth.

Who is this information for?

You have been given this information because your healthcare team think that you may have your baby extremely early (prematurely). You and your family need to know what is likely to happen for you and your baby if this occurs. The maternity team and neonatal (specialist baby doctors and nurses) team will talk to you about this in detail as well as giving you this information and you will have the opportunity to ask any questions that you wish.

What does this mean?

A pregnancy usually lasts for about 40 weeks. How many weeks you are along in your pregnancy (gestation) is usually worked out from an ultrasound scan at around 12 weeks (your dating scan). Babies born before 22 weeks are so small and fragile that they do not survive. Their lungs and other organs are not ready for them to live outside the womb. Many of these babies are not strong enough to survive labour or birth. Even if they do survive labour or birth with signs of life, this will be for a short time and even with the very best neonatal care, they cannot survive for more than a few minutes or hours. Survival focused care is therefore not appropriate for babies born before 22 weeks. Instead, for these babies we will provide care focused on their comfort, and do everything we can to support you to ensure that you are able to spend this precious time making memories with your baby.

Babies born from 22 weeks up to 27 weeks are still incredibly small and fragile. Sometimes babies born this early are not strong enough to survive labour and birth. The earlier the baby is born, the less likely it is that they will be able to survive. If they are born alive, they may be able to survive if they receive intensive medical treatment. However, particularly for babies born below 24 weeks, intensive medical treatment after birth may not be in their best interests, and some extremely premature babies sadly die despite this treatment.

Babies who are born extremely early and survive are also at increased risk of problems with health and development as they grow up. These risks get higher the earlier (more prematurely) a baby is born, and are especially common in those children born before 25 weeks of gestation. Health problems may include breathing difficulties, gut problems (including difficulties with feeding) and eye problems. Developmental problems may include problems with movement, learning and behaviour that can range from mild to very severe; such problems are described on the following page.

The doctors and midwives will talk to you about what they expect for your baby. In some situations, there are difficult decisions to be made about how to care for your baby before and after birth. The right thing to do can be different for different families. That is why it is important that you are fully informed and can discuss with the doctors and midwives what your wishes are for your baby.

Outcome for extremely premature babies born between 22-26 gestation



Chance of baby being born alive

Overall chance of survival
If born alive and receive intensive treatment

Chance of severe disability if survives

22 weeks

1 in 3 die in labour
2 in 3 are liveborn

8 in 10 die
2 in 10 survive



1 in 3 babies has severe disability
2 in 3 do not*

23 weeks

1 in 6 die in labour
5 in 6 are liveborn

6 in 10 die
4 in 10 survive



1 in 4 babies has severe disability
3 in 4 do not*

24 weeks

1 in 10 die in labour
9 in 10 are liveborn

3 in 10 die
7 in 10 survive



1 in 5 babies has severe disability
4 in 5 do not*

25 weeks

1 in 20 die in labour
19 in 20 are liveborn

2 in 10 die
8 in 10 survive




1 in 9 babies has severe disability
8 in 9 do not*

26 weeks

Less than 1 in 20 die in labour
More than 19 in 20 are liveborn

1-2 in 10 die
8-9 in 10 survive



1 in 12 babies has severe disability
11 in 12 do not*

These are the outcomes that are seen on average. For any individual baby, their risk may be higher or lower than this because of other risk factors eg how well grown baby is, where they are born, whether their mother has received steroids etc. Your doctor will tell you if your baby is at higher or lower risk than average.

*Even if your baby is born without a severe disability there is a 1 in 4 chance that they will have other forms of disability such as learning difficulty, cerebral palsy or behavioural problems if they are born before 27 weeks.

‘Outcome’

These pictures are based on what we know about the small number of babies born extremely prematurely in the UK. They show how many babies survive out of every 10 babies born alive this early, and of those who do survive, how many are likely to have a ‘severe disability’ when they grow up.

The majority of babies grow up without severe disability. A proportion of these children will develop other problems as they grow up which may mean, for example, that they need extra help in school or have problems with walking or moving around. Some may have social and emotional problems. The frequency with which children have these problems is greatest the earlier they are born, and problems are most common in children born at 22 to 24 weeks of gestation.

The chance for your baby depends on a number of different things. As well as how early they are born, it also matters how much your baby weighs when it is born, whether it is a boy or girl, whether it is a multiple birth and also how well you and your baby are around the time of birth.

What does ‘severe disability’ mean?

Disability can mean different things to different people. When talking about babies who have been born extremely prematurely, the term severe disability includes problems such as:

- Not being able to walk or even get around independently (this includes conditions such as severe cerebral palsy).
- Being unable to talk, or see or hear properly.
- Difficulties with swallowing or feeding safely.
- Having multiple health problems with frequent visits to hospital.
- Needing to attend separate school for children with special educational needs.
- Being unable to care for themselves or live independently as they grow up.

What does this mean for your baby?

We don’t know exactly the future for your baby. Every baby is different and it is important to talk with your doctors and midwife. They will give you specific information about your own and your baby’s condition.

What can parents do?

What is right for your baby and your family is very individual to you. Your doctors will talk with you about your situation and seek to understand what is important for you and your family. They will support and guide you and involve you in making decisions about treatment for your baby. Thinking about your hopes, your wishes, and your fears about your baby can help the team to support you in the best way possible.

What may happen with my baby?

Not surviving labour and birth: Some babies who are born this early do not survive labour and birth. If this happens you will have the opportunity to spend time with your baby and to make memories with them.

Comfort Care: You and the team may decide that after birth it will be best to provide comfort care to your baby, either because there is an extremely high risk that your baby will not survive or he/she is likely to suffer from life-long disability even with the very best treatment. Comfort care is also known as palliative care and is special care for babies whose time is precious but short. It means providing treatments that will make their time as comfortable as possible. We will help you to be part of this care if you would like. Holding your baby close to you and talking to your baby may be

very comforting.

More information about comfort care or 'palliative care' for babies is available from [Together for Short Lives](#) and [Bliss](#).

Neonatal Intensive Care: You and the team may decide that starting neonatal intensive care would be best for your baby. This will mean you will need some extra treatments before your baby is born. You will be given steroids to help the baby's lungs and brain and magnesium which also helps to protect your baby's brain. You may need to be transferred to a specialist centre, ideally before you have your baby, but there may not be time to do this safely. The team will also talk to you about the treatment that will be given to your baby immediately after birth and what may happen next depending on how your baby reacts to this treatment.

If you and the team decide that intensive care is best for your baby, you should be offered the opportunity to be shown around the neonatal unit (if there is time for this) as it may help to see the neonatal unit and meet the people that work there before your baby is born. You can also talk to staff about expressing breast milk, as this makes such a big difference for premature babies.

What if my baby doesn't come now?

If your baby does not come in the next few days their chances of survival may improve. Ideally, they will stay in the womb for as long as possible as long as you and baby remain healthy.

If that happens there may be different options for you and your baby around the time of birth. That will depend on when your baby comes and on other things that affect the baby's chances of responding to treatment. If this is the case, your healthcare team will continue the conversation with you about what has changed and what different options may be available depending on when your baby is likely to be born, and you will be able to discuss and revise your agreed plans accordingly.

What might my baby look like?

Babies born this early can weigh less than half a kilogram (1 pound or small packet of sugar) and can look quite different to how we imagine a newborn baby. Their skin is shiny and thin and covered with fine hair. Sometimes babies can be quite bruised from the birth. If the baby has died before being born, they will usually be still. Occasionally, where babies have died very close to being born, they may make brief reflex movements that disappear very quickly.

If your baby is born alive, they may take a breath and make a small cry or they may not breathe. Their eyes may not be able to open yet. The baby's colour is often purple or blue to start with.

Transfer to a different hospital

When you have decided with the obstetric and neonatal care teams that starting neonatal intensive care would be best for your baby, research shows that for babies born before 27 weeks of gestation it is best, whenever possible, to be born in a specialist maternity unit with a specialist Neonatal Intensive Care Unit (sometimes called a 'Level 3 NICU'). If a baby born before 27 weeks of gestation is born in a maternity unit (or at home) where there is not a specialist NICU, then we know that the baby will generally do better if moved to a specialist NICU after birth.

If your hospital does not have a specialist NICU, this may mean that you will be offered transfer to one of these centres before your baby is born. We understand that this can be a very anxious time and that you may be moved quite some distance from home. It can be very difficult to predict which mothers will deliver early and so some mothers may be moved to another hospital and their baby not born early.

It may also be the case that you are considered too unwell or too far on in labour to be safely moved to another hospital before your baby is born. When it is not possible to transfer you before the baby has been born your baby may be transferred by a specialist Neonatal Transport Team after the birth. Your own health needs may mean you will be unable to travel immediately with your baby but your local maternity team will do everything they can to move you to the same unit as your baby as soon as it is safe to do so.

When your baby is well and stable and no longer requiring neonatal intensive care, the team will discuss the transfer of your baby to a neonatal unit closer to your home (repatriation)

We appreciate that moving to another hospital can be distressing for you and your family, especially if you are separated from your baby for a while. We will talk to you about this in more detail if it is decided that this is the best option for your family.

What if I have more questions?

This information has been provided to you as part of the conversation that your healthcare team will have with you about your baby. If you have any other questions do make sure you ask your doctors and nurses to answer them, so you have all the information you need about your situation and the options available to you. Your healthcare team want to work with you make the best decision for your baby and for your family.

This space is for the health care team who are discussing this with you to write extra details about your baby or babies.

You may want to use this space to write down some questions to discuss with the team.

Many families find it useful to have follow-up discussions, so please ask to speak to the neonatal and maternity team again at any point. We know this is a lot of information to take in, and many families find it useful to have further discussions with the healthcare team once they have had a chance to think everything through. Your healthcare team will arrange a time to have a follow-up conversation with you where you can ask more questions and discuss this further.

Useful contact details:

Bliss - Premature and sick baby charity
www.bliss.org.uk

Together for Short Lives - Charity for babies and children with life-limiting conditions
www.togetherforshortlives.org.uk
Helpline: 0808 8088 100

Sands - Stillbirth and neonatal death charity
www.uk-sands.org
Helpline: 0808 1643332
Email helpline@sands.org.uk

References

Backes CH, Rivera BK, Pavlek L, Beer LJ, Ball MK, Zettler ET, Smith CV, Bridge JA, Bell EF, Frey HA. Proactive neonatal treatment at 22 weeks of gestation: a systematic review and meta-analysis. *Am J Obstet Gynecol*. 2021;224(2):158-174.

British Association of Perinatal Medicine (BAPM) Framework for Practice: Lactation and Loss. Available at <https://www.bapm.org/resources/lactation-and-loss-management-of-lactation-following-the-death-of-a-baby> Accessed 7/5/25

BAPM Framework for Practice: Pre-hospital management of the baby born at extreme preterm gestation. Available at <https://www.bapm.org/resources/pre-hospital-management-of-the-baby-born-at-extreme-preterm-gestation> Accessed 7/5/25

BAPM Framework for Practice: Recognising uncertainty: an integrated framework for palliative care in perinatal medicine. Available at <https://www.bapm.org/resources/palliative-care-in-perinatal-medicine-framework> Accessed 7/5/25

BAPM Nanopreterm Special Interest Group. Available at <https://www.bapm.org/pages/nano-preterm> accessed 31/7/25

BAPM Neonatal Airway Safety Standard. Available at <https://www.bapm.org/resources/BAPM-Neonatal-Airway-Safety-Standard> Accessed 7/5/25

BAPM Perinatal Optimisation Pathway. Available at <https://www.bapm.org/pages/perinatal-optimisation-pathway> Accessed 7/5/25

BAPM Perinatal Optimisation Resources. Available at <https://www.bapm.org/pages/perinatal-optimisation-pathway-resources> Accessed 7/5/25

BAPM Practice Guide: In-utero Transfer. Available at <https://www.bapm.org/resources/in-utero-transfer> Accessed 13/5/25

Bell EF, Hintz SR, Hansen NI, Bann CM, Wyckoff MH, DeMauro SB, Walsh MC, Vohr BR, Stoll BJ, Carlo WA, Van Meurs KP, Rysavy MA, Patel RM, Merhar SL, Sánchez PJ, Laptook AR, Hibbs AM, Cotten CM, D'Angio CT, Winter S, Fuller J, Das A; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Mortality, In-Hospital Morbidity, Care Practices, and 2-Year Outcomes for Extremely Preterm Infants in the US, 2013-2018. *JAMA*. 2022;327(3):248-263.

Blanc J, Resseguier N, Goffinet F, Lortie E, Kayem G, Delorme P, Vayssi re C, Auquier P, D'Ercole C. Association between gestational age and severe maternal morbidity and mortality of preterm caesarean delivery: a population-based cohort study. *Am J Obstet Gynecol* 2019;220:399.e1-9. Bliss Baby Charter. Available at <https://www.bliss.org.uk/health-professionals/bliss-baby-charter> accessed 11/5/25

Bliss and Together for Short Lives. Information for Parents Document: 'Caring for your baby when the future is uncertain' 2024. Available at <https://www.bliss.org.uk/parents/uncertainty-and-loss> accessed 31/7/25

Costeloe K, Hennessy E, Gibson AT, Marlow N, Wilkinson AR. The EPICure study: Outcome to

discharge from hospital for infants at the threshold of viability. *Pediatrics* 2000;106:659-671.

David AL, Soe A. Extreme prematurity and perinatal management. *TOG* 2018;20:109-17.

Doyle LW, Crowther CA, Middleton P, Marret S, Rouse D. Magnesium sulphate for women at risk of preterm birth for neuroprotection of the fetus. *Cochrane Database of Systematic Reviews* 2009, Issue 1. Art. No.: CD004661. DOI: 10.1002/14651858.CD004661.pub3

Edwards EM, Ehret DEY, Soll RF, Horbar Survival of Infants Born at 22 to 25 Weeks' Gestation Receiving Care in the NICU: 2020-2022. *Pediatrics*. 2024;154(4):e2024065963.

Ehret DEY, Edwards EM, Greenberg LT, Bernstein IM, Buzas JS, Soll RF, Horbar JD. Association of antenatal steroid exposure with survival among infants receiving postnatal life support at 22 to 25 weeks' gestation. *JAMA Network Open*. 2018;1(6):e183235.

Farooqi A, Hakansson S, Serenius F, Kallen K, Björklund L, Normann E, Domellöf M, Ådén U, Abrahamsson T, Elfvin A, Sävman K, Bergström PU, Stephansson O, Ley D, Hellstrom-Westas L, Norman M. One-year survival and outcomes of infants born at 22 and 23 weeks of gestation in Sweden 2004-2007, 2014-2016 and 2017-2019. *Arch Dis Child Fetal Neonatal Ed*. 2023;109(1):10-17.

Fogarty M, Osborn DA, Askie L, Seidler AL, Hunter K, Lui K, Simes J, Tarnow-Mordi W. Delayed vs early umbilical cord clamping for preterm infants: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2018;218:1-8

Gallagher K, Marlow N, Edgley A, Porock D. The attitudes of neonatal nurses towards extremely preterm infants. *J. Adv. Nurs*. 2012;68:1768-79.

Gallagher K, Aladangady N, Marlow N. The attitudes of neonatologists towards extremely preterm infants: a Q methodological study. *Arch Dis Chil Fetal Neonatal Ed* 2016;101:31-36.

Goodfellow L, Care A, Curran C, Roberts D, Turner MA, Knight M, Zarko A. Preterm prelabour rupture of membranes before 23 weeks' gestation: prospective observational study: *BMJ Medicine* 2024;3:e000729.

Grabovac M, Karim JN, Isayama T, Korale Liyanage S, McDonald SD. What is the safest mode of birth for extremely preterm breech singleton infants who are actively resuscitated? A systematic review and meta-analyses. *BJOG* 2018;125:652-663.

Griffin M, Smith-Collins A, Hodgetts-Morton V, Hughes T, Magill L, Perry R, Morris K on behalf of the TRANSFER project collaborative, UK. TRANSFER:Threatened preterm birth, assessment of the need for in-utero transfer between 22+0-23+6 weeks' gestation, a multicentre prospective service evaluation across the United Kingdom. *Eur J Obstet Gynecol Reprod Biol* 2025;309:73-78

Guillén Ú, Weiss EM, Munson D, Maton P, Jefferies A, Norman M, Naulaers G, Mendes J, Justo da Silva L, Zoben P, Hansen TW, Hallman M, Delivoria-Papadopoulos M, Hosono S, Albersheim SG, Williams C, Boyle E, Lui K, Darlow B, Kirpalani H. Guidelines for the Management of Extremely Premature Deliveries: A Systematic Review. *Pediatrics* 2015;136:343-50.

Haines M, Wright IM, Bajuk B, Abdel-Latif ME, Hilder L, Challis D, Guaran R, Oei JL. Population-based study shows that resuscitating apparently stillborn extremely preterm babies is associated with poor outcomes. *Acta Paediatr* 2016;105:1305-11.

Helenius K, Longford N, Lehtonen L, Modi N, Gale, C on behalf of the Neonatal Data Analysis Unit and the United Kingdom Neonatal Collaborative. Association of early postnatal transfer and birth outside a tertiary hospital with mortality and severe brain injury in extremely preterm infants: observational cohort study with propensity matching. *BMJ* 2019; 367:15678

Joint Royal Colleges Ambulance Liaison Committee, Association of Ambulance Chief Executives (2021) JRCALC Clinical Guidelines. Cited from: JRCALC Plus (2021) [Mobile application software]. Bridgwater: Class Publishing Ltd.

Kaempf JW, Tomlinson MW, Campbell B, Ferguson L, Stewart VT. Counseling pregnant women who may deliver extremely premature infants: medical care guidelines, family choices, and neonatal outcomes. *Pediatrics* 2009;123:1509-15.

Kharrat A, Moore GP, Beckett S, Nicholls SG, Sampson M, Daboval T. Antenatal consultations at extreme prematurity: a systematic review of parent communication needs. *J Pediatr* 2018;196:109-15.

Kono Y, Yonemoto N, Nakanishi H, Kusuda S, Fujimura M. Changes in survival and neurodevelopmental outcomes of infants born at <25 weeks gestation: a retrospective observational study in tertiary centres in Japan. *BMJ Paediatrics Open* 2018;2:e000211.

Larcher V, Craig F, Bhogal K, Wilkinson D, Brierley J, on behalf of the Royal College of Paediatrics and Child Health. Making decisions to limit treatment in life-limiting and life-threatening conditions in children: a framework for practice. *Arch Dis Child* 2015;100:s1-s23.

LeMoine FV, Battarbee AN, Travers CP, Battersby C, Gibson KS; Tiny Baby Collaborative Steering Committee. Considerations for obstetric management of births 22-25 weeks' gestation. *J Perinatol*. 2025 Apr 11. doi: 10.1038/s41372-025-02289-y. Epub ahead of print.

Lemyre B, Moore G. Counselling and management for anticipated extremely preterm birth. *J Paediatr Child Health* 2017;22:334–341.

Li YX, Hu YL, Huang X, Li J, Li X, Shi ZY, Yang R, Zhang X, Li Y, Chen Q. Survival outcomes among periviable infants: a systematic review and meta-analysis comparing different income countries and time periods. *Front Public Health*. 2024;12:1454433.

Macfarlane PI, Wood S, Bennett J. Non-viable delivery at 20-23 weeks gestation: observations and signs of life after birth. *Arch Dis Child Fetal Neonatal Ed* 2003;88: F199-202.

Mactier H, Bates SE, Johnston T, Lee-Davey C, Marlow N, Mulley K, Smith LK, To M, Wilkinson D, BAPM Working Group. Perinatal management of extreme preterm birth before 27 weeks of gestation: a framework for practice. *Arch Dis Child Fetal Neonatal Ed* 2020;105:232-239.

Manley BJ, Dawson JA, Kamlin CO, Donath SM, Morley CJ, Davis PG. Clinical assessment of extremely premature infants in the delivery room is a poor predictor of survival. *Pediatrics* 2010;125:559-64.

Marlow N, Wolke D, Bracewell M, Samara M, for the EPICure Study Group. Neurologic and developmental disability at 6 years of age after extremely preterm birth. *N Engl J Med* 2005; 352:9-19.

Marlow N, Bennett C, Draper ES, Hennessy EM, Morgan AS, Costeloe KL. Perinatal outcomes for extremely preterm babies in relation to place of birth in England: the EPICure 2 study. *Arch Dis Child*

Fetal Neonatal Ed 2014;99:F181–F188.

McGrath JS, Roehr CC, Wilkinson D. When should resuscitation at birth cease? *Early Hum Dev* 2016; 101:31-36

Mehler K, Oberthuer A, Keller T, Becker I, Valter M, Roth B, Kribs A. Survival among infants born at 22 or 23 weeks' gestation following active prenatal and postnatal care. *JAMA Pediatrics* 2016;170:671–77.

Mother and Babies: Reducing Risk through Audits and Confidential Enquiries across the UK (MBRRACE-UK) Guidance 2020: Determination of signs of life following spontaneous birth before 24⁺⁰ weeks of gestation where, following discussion with the parents, active survival-focused care is not appropriate. Available at <https://timms.le.ac.uk/signs-of-life/guidance.html> Accessed 7/5/25

Myrhaug HT, Brurberg KG, Hov L, Markestad T. Survival and impairment of extremely premature infants: a meta-analysis. *Pediatrics* 2019;143(2):e20180933

National Institute for Clinical Excellence (NICE) Guidelines NG25 Preterm Labour and Birth, updated 2022. Available at <https://www.nice.org.uk/guidance/ng25/chapter/Update-information> accessed 8/9/19

NICE Guidelines NG201 Antenatal Care for Uncomplicated Pregnancies, updated 2021. Available at <https://www.nice.org.uk/guidance/ng201> accessed 13/5/25

National Bereavement Care (NBC) Pathway in England Available at: <https://www.nbcpathway.org.uk/pathways/neonatal-death-bereavement-care-pathway/> accessed 11/5/25

National Bereavement Care (NBC) Pathway in Scotland Available at: https://www.nbcscotland.org.uk/media/vlymy5hc/nbc-scotland-neonatal-death-pathway_03.pdf accessed 31/7/25

National Bereavement Care (NBC) Pathway for Termination of Pregnancy for Fetal Anomaly (ToPFA) 2022. Available at: <https://www.nbcpathway.org.uk/pathways/termination-of-pregnancy-for-fetal-anomaly-topfa-bereavement-care-pathway/> accessed 31/7/25

National Neonatal Audit Programme (NNAP) Extended Analysis Report 2023 data. Page 21, Table 5. Available at: https://www.rcpch.ac.uk/sites/default/files/2024-10/nnap_2023_data_extended_analysis_report_v2.pdf accessed 31/7/25

Neonatal Critical Care Review (NCCR) <https://www.england.nhs.uk/wp-content/uploads/2019/12/Implementing-the-Recommendations-of-the-Neonatal-Critical-Care-Transformation-Review-FINAL.pdf> accessed 31/7/25

Neonatal Service Quality Indicators (NSQI) Standards relating to Structures and Processes supporting Quality and Patient Safety in Neonatal Services. BAPM, June 2017. Available at https://hubble-live-assets.s3.eu-west-1.amazonaws.com/bapm/file_asset/file/86/NSQI_FINAL_0.pdf accessed 11/5/25

NHS England Saving Babies Lives Care Bundle (version 3) <https://www.england.nhs.uk/publication/saving-babies-lives-version-three/> accessed 13/5/25

NHS England Medical Examiners System <https://www.england.nhs.uk/patient-safety/medical-examiners/the-national-medical-examiner-system/> Accessed 13/5/25

Norman M, Hallberg B, Abrahamsson T, Björklund LJ, Domellöf M, Farooqi A, Foyen Bruun C, Gadsbøll C, Hellström-Westas L, Ingemansson F, Källén K, Ley D, Maršál K, Normann E, Serenius F, Stephansson O, Stigson L, Um-Bergström P, Håkansson S. Association Between Year of Birth and 1-Year Survival Among Extremely Preterm Infants in Sweden During 2004-2007 and 2014-2016. *JAMA*. 2019 Mar 26;321(12):1188-1199.

Nuffield Council on Bioethics. Critical care decisions in fetal and neonatal medicine: ethical issues. 2006. www.nuffieldbioethics.org

Online Confidence Interval calculator. Kohn MA, Senyak J. Sample Size Calculators [website]. UCSF CTSI. 28 September 2024. Available at <https://www.sample-size.net/> Accessed 26/2/25.

Papageorgiou AT, Kemp B, Stones W, Ohuma EO, Kennedy SH, Purwar M, Salomon LJ, Altman DG, Noble JA, Bertino E, Gravett MG, Pang R, Cheikh Ismail L, Barros FC, Lambert A, Jaffer YA, Victora CG, Bhutta ZA, Villar J; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). Ultrasound-based gestational-age estimation in late pregnancy. *Ultrasound Obstet Gynecol* 2016;48:719-726.

Park GY, Park WS, Yoo HS, Ahn SY, Sung SI, Kim SS, Chang YS. Short-term outcomes comparison between preterm infants with and without acute hypoxic respiratory failure attributable to presumed pulmonary hypoplasia after prolonged preterm premature rupture of membranes before 25 gestational weeks. *J Matern Fetal Neonatal Med* 2018;13:1-8.

Patel RM, Rysavy MA, Bell EF, Tyson JE. Survival of Infants Born at Perivable Gestational Ages. *Clin Perinatol* 2017;44:287–303.

Pearl S, Ray O, Galletta L, Bates A, Boland RA, Davis PG, Gale C, Johnson S, Kinsella S, Knight M, Owen LS, Pallot L, Prentice TM, Santhanadass P, Stanbury K, Tingay D, Whitehead CL, Manley BJ, Roehr CC, Hardy P; Steering Group. Research priorities for the most premature babies born <25 weeks' gestation: results of an international priority setting partnership. *Arch Dis Child Fetal Neonatal Ed*. Published Online First: 23 February 2025. doi: 10.1136/archdischild-2024-328133

POPPY Steering Group, Family-centred care in neonatal units: A summary of research results and recommendations from the POPPY Project, 2009; NHS England, Neonatal Critical Care Transformation Review, 2018

Reddy UM, Rice MM, Grobman WA, Bailit JL, Wapner RJ, Varner MW, Thorp JM Jr, Leveno KJ, Caritis SN, Prasad M, Tita AT, Saade GR, Sorokin Y, Rouse DJ, Blackwell SC, Tolosa JE; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Serious maternal complications after early preterm delivery (24– 33 weeks' gestation). *Am J Obstet Gynecol* 2015;213:538.e1-9.

Resuscitation Council <https://www.resus.org.uk/library/2021-resuscitation-guidelines/newborn-resuscitation-and-support-transition-infants-birth> Accessed 11/5/25

Roberts D, Brown J, Medley N Dalziel SR. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database of Systematic Reviews* 2017, Issue 3. Art. No.: CD004454. DOI: 10.1002/14651858.CD004454.pub3

Royal College of Obstetricians and Gynaecologists (RCOG) Green Top Guideline (GTG) No. 36, 2017.

Prevention of early-onset neonatal group B streptococcal disease. Available at <https://www.rcog.org.uk/guidance/browse-all-guidance/green-top-guidelines/prevention-of-early-onset-group-b-streptococcal-disease-green-top-guideline-no-36/> accessed 31/7/25

Royal College of Obstetricians and Gynaecologists (RCOG) Green Top Guideline (GTG) No. 73, 2019. Care of women presenting with suspected preterm prelabour rupture of membranes from 24+0 weeks of gestation. Available at <https://www.rcog.org.uk/guidance/browse-all-guidance/green-top-guidelines/care-of-women-presenting-with-suspected-preterm-prelabour-rupture-of-membranes-from-24plus0-weeks-of-gestation-green-top-guideline-no-73/> accessed 31/7/25

Royal College of Obstetricians and Gynaecologists (RCOG) Position Statement following Chief Coroner's Guidance No.45 "Stillbirth and Live Birth Following Termination of Pregnancy" December 2023. Available at: <https://www.rcog.org.uk/media/ny1pc5ml/position-statement-coroners-guidance-no-45.pdf> accessed 12/5/25

Royal College of Physicians and Surgeons of Glasgow (RCPSG) Policy Statement <https://rcpsg.ac.uk/college/this-is-what-we-stand-for/policy/consent/the-montgomery-case> Accessed 4/9/19

Royal College of Paediatrics and Child Health (RCPCH) 2019 <https://www.rcpch.ac.uk/news-events/news/new-advice-seeks-prevent-parent-doctor-conflict-paediatrics> accessed 7/5/25

Rysavy MA, Li L, Bell EF, Das A, Hintz SR, Stoll BJ, Vohr BR, Carlo WA, Shankaran S, Walsh MC, Tyson JE, Cotten CM, Smith PB, Murray JC, Colaizy TT, Brumbaugh JE, Higgins RD; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Between-hospital variation in treatment and outcomes in extremely preterm infants. *N Engl J Med* 2015;372:1801-11.

Serenius F1, Källén K, Blennow M, Ewald U, Fellman V, Holmström G, Lindberg E, Lundqvist P, Maršál K, Norman M, Olhager E, Stigson L, Stjernqvist K, Vollmer B, Strömberg B; EXPRESS Group. Neurodevelopmental outcome in extremely preterm infants at 2.5 years after active perinatal care in Sweden. *JAMA*. 2013 May 1;309(17):1810-20.

Smith LK, van Blankenstein E, Fox G, Seaton SE, Martínez-Jiménez M, Petrou S, Battersby C; MBRRACE-UK Perinatal Surveillance Group. Effect of national guidance on survival for babies born at 22 weeks gestation in England and Wales: population based cohort study. *BMJ Medicine* 2023;2:e000579.

Together for Short Lives: <https://www.togetherforshortlives.org.uk/wp-content/uploads/2018/01/ProRes-Core-Care-Pathway.pdf> Accessed 3/5/19

Travers CP, Clark RH, Spitzer AR, Das Am Garite TJ, Carlo WA. Exposure to any antenatal corticosteroids and outcomes in preterm infants by gestational age: prospective cohort study. *BMJ* 2017;356:j1039

Van Blankenstein E, Sodiwala T, Lanoue J, Modi N, Uthaya SN, Battersby C; UK Neonatal Collaborative. Two-year neurodevelopmental data for preterm infants born over an 11-year period in England and Wales, 2008–2018: a retrospective study using the National Neonatal Research Database. *Arch Dis Child Fetal Neonatal Ed* 2024;109:143-150.

Wilkinson D, Verhagen E. Thresholds for resuscitation of extremely preterm infants in the UK, Sweden, and Netherlands. *Pediatrics* 2018;142:s1

Wilkinson D. Death or Disability? The Carmentis machine and decision making for critically ill children. Oxford University Press. 2013

Wilkinson D, Marlow N, Hayden D, Mactier H. Recommendations in the face of uncertainty: should extremely preterm infants receive chest compressions and/or epinephrine in the delivery room? *Arch Dis Child Fetal Neonatal Ed.* 2020;105(3):240-241.

Wyckoff HM, Salhab WA, Heyne RJ, Kendrick DE, Stoll BJ, Laptook AR. Outcome of extremely low birth weight infants who received delivery room cardiopulmonary resuscitation. *J Pediatr* 2012;160:239-44.

Young E, Tsai E, O’Riordan A. A qualitative study of predelivery counselling for extreme prematurity. *Paediatr Child Health* 2012;17:432-6.

Xiao D, Zhu T, Qu Y, Gou X, Huang Q, Li X, Mu D. Maternal chorioamnionitis and neurodevelopmental outcomes in preterm and very preterm neonates: A meta-analysis. *PLoS One* 2018;13(12); (12):e0208302



BAPM

Leading Excellence in Perinatal Care

**This document was produced by the
British Association of Perinatal Medicine (BAPM).**

BAPM is a membership organisation that is here to support all those involved in perinatal care to optimise their skills and knowledge, deliver and share high-quality safe and innovative practice, undertake research, and speak out for babies and their families.

We are a professional association of neonatologists, paediatricians, obstetricians, nurses, midwives, trainees, network managers and other health professionals dedicated to shaping the delivery and improving the standard of perinatal care in the UK.

Our vision is for every baby and their family to receive the highest standard of perinatal care. Join us today.

www.bapm.org/join

British Association of Perinatal Medicine (BAPM)
is registered in England & Wales
under charity number 1199712 at
5-11 Theobalds Road, London, WC1X 8SH