



Review

Parental health in the context of public family care proceedings: A scoping review of evidence and interventions

Claire Grant^{a,*}, Jessica Radley^b, Georgia Philip^c, Rebecca Lacey^a, Ruth Blackburn^d,
Claire Powell^d, Jenny Woodman^e

^a Department of Epidemiology and Public Health, University College London, UK

^b Institute of Psychiatry, Psychology and Neuroscience, Kings College London, UK

^c Centre for Research on Children & Families, University of East Anglia, Norwich, UK

^d Great Ormond Street Institute of Child Health, University College London, London, UK

^e Thomas Coram Research Unit, Social Research Institute, University College London, UK

ARTICLE INFO

Keywords:

Child protection

Parental health

Social work

Family studies

ABSTRACT

Background: Child protective services (CPS), or their equivalent, have statutory power to remove children from birth parents in instances of child abuse, neglect, or concerns around parenting capacity via public family care proceedings. Parents who have children subject to proceedings, ‘birth parents’, often have complex health and social care needs.

Objective: We aimed to review what is known about the health needs of birth parents and the interventions implemented to support these health needs.

Methods: We searched PubMed, Scopus, and grey literature using a systematic strategy of key concepts “health”, “care proceedings”, and “parents”. We included all publications in English that reported parental health in the context of care proceedings from the 1st of January 2000 to the 1st of March 2021.

Results: Included studies ($n = 61$) reported on maternal health (57 %) or the health of both parents (40 %), with only one study reporting on fathers alone. We conceptually categorised parental health need ($n = 41$) into i) mental health, ii) physical health, iii) substance misuse, iv) developmental disorders, and v) reproductive health. Health inequities and poor access to services were described across all categories, with longstanding issues often pre-dating proceedings or the child’s birth. All interventions supporting parental health ($n = 20$) were targeted at mothers, with some supporting fathers ($n = 8$), formally or informally. We grouped similar interventions into three types: alternative family courts, wrap-around services, and specialist advocacy/peer support.

Conclusions: Parents who have children subject to care proceedings have complex health needs that pre-date CPS concerns. The studies included in our review strongly suggest that health issues are exacerbated by child removal, triggering deteriorations in mental health, poor antenatal health for subsequent pregnancies, and avoidable mortality. Findings highlight the need for targeted and timely intervention for parents to improve whole-family outcomes. There are models that have been designed, implemented, and tested using relationship-based, trauma-informed, multidisciplinary, family-focused, and long-term approaches.

* Corresponding author.

E-mail address: claire.grant.20@ucl.ac.uk (C. Grant).




Fig. 1. Flow diagram for inclusion as outlined in PRISMA statement.

1. Background

Underpinned by the United Nation's Convention on the Rights of the Child agreement, governments across the world have a duty to protect children (United Nations, 1989). Child protective services (CPS), or equivalent systems, provide a method for a state to intervene in a child's care when there are significant concerns over their welfare. Such public 'care proceedings' provide statutory power to remove a child from their birth parents when necessary (Gilbert et al., 2011). Most commonly, this is in instances of child maltreatment (abuse, neglect) or concerns around parenting capacity (UK Public General Acts, 1989). States have responsibility to protect at-risk children, yet there are difficulties in balancing the harms and benefits of such intrusive interventions into family life (Munro & Ward, 2008). The decision for a child to be taken into care has enduring consequences for the child and the parent, with Looked After children experiencing poorer health, social and education outcomes compared to that of their peers (Berlin et al., 2011; Courtney et al., 2007). Looked After children are also more than twice as likely than other adolescents to enter parenthood early and to have CPS involvement and intervention with their own children (Coman & Devaney, 2011; Wall-Wieler, Roos, Nickel, Chateau, and Brownell, 2018f).

The biological parents of children subject to care proceedings ('birth parents') often have complex health needs, such as mental health and substance misuse difficulties, which can lead to CPS involvement and intervention with their children (Bedston et al., 2019; Broadhurst et al., 2017; Philip et al., 2021). Many people with complex health needs care adequately for their children, yet some health challenges can directly and indirectly impact on parenting ability (Barlow et al., 2006; Munro & Ward, 2008). For example, studies have reported increased risk of child maltreatment and accidental injury among families with parental substance misuse and mental health difficulties (Nevriana et al., 2020; Pierce et al., 2020). Parents may also experience periods of relapse or acute illness requiring hospitalisation, affecting their ability to take care of children and family functioning (Källquist & Salzmänn-Erikson, 2019).

There is increasing awareness of the interrelated health needs of children and their caregivers (Woodman et al., 2020). Although it is not always possible to prevent children being Looked After, public services have an opportunity to support parental health to interrupt lifelong and intergenerational disadvantage (Bywaters et al., 2016; UK Government, 2022). Given that care proceedings themselves are likely to worsen health issues due to heightened stress and threat to parental identity, there is also an ethical imperative to help these parents (Broadhurst & Mason, 2013; Family Rights Group, 2018). Targeted and effective intervention could result in fewer children being removed from their families, including any potential future pregnancies and subsequent care proceedings (Broadhurst, Alrouh, Yeend, Harwin, Shaw, Pilling, Mason, and Kershaw, 2015a; Skinner et al., 2021). The first step in policy and practice change to support birth parents is to understand their health needs and the interventions and practice approaches that feasibly and effectively address these needs. We understand many health inequalities are likely downstream effects of entrenched social and economic inequalities; however, this is outside the scope of this review.

1.1. Study objectives

We reviewed the existing literature on the health needs of birth parents before, during and after care proceedings and interventions or practices which had been evaluated in terms of addressing the health needs of birth parents.

2. Methods

We conducted a scoping review using a systematic approach to enhance robustness (Munn et al., 2018). We report our results based on PRISMA-ESR (Tricco et al., 2018). Our protocol has been published (Grant et al., 2021).

2.1. Eligibility criteria

We included all original research published in English since 2000 that reported on the health of parents whose children were subject to care proceedings. Care proceedings were defined as the involvement of public services in determining child placement for at-risk children under 18. Parental health need was conceptualised as physical and/or mental health needs that could benefit from services, such as health education, disease prevention, diagnosis, treatment, or rehabilitation (Public Health England, 2014). All reviews, opinion pieces or descriptions of health interventions without evaluations were excluded, but reference lists were screened to check for relevant literature (see supplementary material 1).

2.2. Searches

We systematically searched two scientific databases (PubMed and Scopus) and grey literature sources using key concepts 'health need', 'care proceedings' and 'parents' (see Fig. 1). All titles and abstracts returned were screened by two independent reviewers (CG & JR) with a 91 % agreement rate. In instances of disagreement, the title/abstract was revisited, and consensus reached. CG conducted all full-text screening (see supplementary material 2).

2.3. Data extraction and synthesis

We extracted information on health need, methods, and results for all studies. For included interventions, information on health

Table 1
Parental health need summary.

Health need (n = studies)	Countries (n = studies)	Inclusive of fathers (% of sample)	Summary findings for birth parents	
			Qualitative findings	Quantitative findings
Mental health (n = 20)	UK (n = 7) USA (n = 5) Australia (n = 4) Canada (n = 2) Germany (n = 1) Denmark (n = 1)	✓ 12 % of qualitative <1 % of quantitative	- Adverse childhood experiences and trauma common among birth parents and often associated with complex mental health need - Mental health problems raised as professional concern for parenting capacity in care proceedings - Identity of being a 'parent' as an important motivator for recovery and meaning among parents with mental illness - Complex feelings of loss and grief following child removal, including a renegotiation of parental identity, and worsening mental health.	- High rates of diagnosis of serious mental illness (SMIs) (schizophrenia spectrum disorders & personality disorders) among birth mothers - Diagnosis of SMI and histories of inpatient psychiatric care as risk factor for child removal, an effect greater for birth mothers - Child removal associated with deteriorating parental mental health, including increased psychiatric prescriptions, rates of anxiety/depression, and suicide attempts and completions.
Physical health (n = 7)	UK (n = 5) Canada (n = 2)	✓ 100 % of qualitative 44 % of quantitative	- Birth fathers reported chronic and long-standing physical health conditions - Physical health concerns not frequently captured in court case files.	- Birth parents are dying earlier than comparator groups and from preventable/amenable causes - Maternal non-engagement with GP associated with child removal - Associations between child removal and self-reported poor health among birth mothers.
Substance misuse (n = 15)	UK (n = 7) USA (n = 5) Australia (n = 1) Finland (n = 1) Canada (n = 1)	✓ 25 % of qualitative 1 % of quantitative	- Substance use perceived as a coping strategy, even before entry to parenthood - Substances used included alcohol, marijuana, methamphetamines, heroin, and opioid-based prescription medication - Substance use raised as professional concern for parenting capacity in care proceedings - Child removal triggering worsening substance misuse, as parents 'self-medicate' acute grief - Co-dependency of both parents using was felt to create unhealthy and destructive environments.	- High rates of substance misuse recorded in birth parents' administrative health and court records - Parental drug use as risk factor for child removal, an effect greater for birth mothers - Risk of child removal greatest for mothers using throughout pregnancy, poly drug users and women living with other drug users.
IDD (N = 14)	UK (n = 7) USA (n = 2) Australia (n = 2) Canada (n = 1) Norway (n = 1) Iceland (n = 1)	✓ 24 % of qualitative 7 % of quantitative	- Feelings of powerlessness for women with IDDs 'mothering differently' - Confusion around court processes and legal jargon used throughout proceedings - IDDs used as proxy for parenting incapacity in court records, with discriminatory evidence used in court [e.g., IQ scores] - Professionals feeling ill-informed at how to support families with IDD	- IDDs included learning, cognitive and behaviour difficulties, ASD, and ADHD - Parents with IDDs overrepresented in care proceedings and child removal orders - Rates of early (newborn or infant) child removal far greater among parents with IDDs - Most at risk were mothers with co-morbid mental illnesses, single mothers and women who received poor antenatal care.
Reproductive health (n = 5)	UK (n = 3) Canada (n = 2)	No fathers		- Birth mothers enter parenthood early - Poor antenatal care associated with increased risk of child removal - Histories of child removal associated with poor antenatal care for subsequent pregnancies - Maternal engagement with primary care contraception services reduced likelihood of child removal.

focus, approach and effectiveness were extracted. We synthesised study findings and included individually reported odds ratios (OR), risk ratios (RR) and hazard ratios (HR) related to parental health outcomes. Statistical results demonstrated the probability that birth parents would experience a particular outcome (OR/RR) or how often particular outcomes happened over time (HR) compared to other groups. We conducted a narrative synthesis of all results with comment and input from a panel of mothers who had lived experience of child removal in England (see supplementary material 3).

3. Results

We included 61 studies reporting on both parental health need ($n = 41$) and supporting interventions ($n = 20$) (see supplementary material 4).

3.1. Study characteristics and populations

Our review captured a huge range of birth parents from across the globe. In the UK, Canada, Australia, and Sweden, administrative health records of over 27,000 birth mothers, 3690 birth fathers and 1280 children with parental information recorded were reported. Such approaches allow large populations to be analysed and can ascertain temporality of events yet cannot determine health need that is not known to health services. Further evidence on parental health was captured in analyses of around 1500 family court case files across the UK, Iceland, Germany, USA, and Australia. Comparison of these data to parents who are not under CPS scrutiny is challenging, as records focus on conditions that impact most heavily on parenting capacity (i.e., complex unmet health needs). Longitudinal data from Australia, Denmark, and the USA reported on the health needs of over 42,840 families involved in care proceedings, although there are issues with sampling biases. Parental health experiences are explored in qualitative interview data from over 190 birth mothers, 25 birth fathers, and 15 allied professionals in the UK, USA, and Australia. These accounts provide a nuanced insight into parental health need, however, can be limited in reflecting service contact.

3.2. Parental health need ($n = 41$)

Studies reporting on parental health need captured mental health ($n = 20$), physical health ($n = 7$), substance misuse and addictions ($n = 15$), intellectual and developmental disabilities ($n = 14$), and reproductive health ($n = 5$) (see Table 1). Most of these studies focused on maternal health ($n = 22$) or both parents ($n = 18$), with only one reporting on fathers exclusively.

3.2.1. Mental health ($n = 20$)

Findings report high rates of mental health need and service use among birth parents, a finding consistent across country, study design and measure. Studies reported high rates of maternal mental health difficulties prior to CPS involvement, including histories of specialist service contact (Griffiths et al., 2020a, b; Pearson et al., 2020a, b, 2021; Salzer et al., 2020; Simkiss et al., 2012; Vigod et al., 2018). Compared to women accessing similar services, birth mothers had higher rates of being diagnosed with serious mental illnesses (SMIs), (personality disorders 21 % vs 11 %, schizophrenia spectrum disorders 19 % vs 11 %) and being admitted to inpatient stay (27 % vs 14 %) (Pearson et al., 2021). Recording of maternal mental illness in GP data was associated with child removal (OR 2.51, 95 % CI 1.55–4.05) (Simkiss et al., 2012) and at a local authority level, maternal adversity (including mental health) accounted for 24 % variation in child removal rates (Pearson et al., 2020a, b).

Mothers with SMIs were more likely to have children placed in out-of-home care than other mothers with CPS involvement in the USA (OR 2.8, 95 % CI = 1.5–5.2) and Canada (OR 6.69, 95 % CI = 3.89–11.52) (Hollingsworth, 2004; Park et al., 2006; Wall-Wieler, Roos, Brownell, Nickel, Chateau, and Nixon, 2018d). Mental health disorders were common professional concerns for parenting capacity in care proceeding files (Broadhurst et al., 2017; Kohl et al., 2011; Kratky & Schröder-Abé, 2018; Sheehan & Levine, 2005), with mothers' poor mental health a greater risk factors for out-of-home child placement (OR 2.33, 95 % CI 2.05–2.63) than fathers (OR 1.06, CI 95 % 0.94–1.19) (Whitten et al., 2021). The risk of custody loss was greatest for women with pre-existing mental illness (OR 4.77, 95 % CI 4.13–5.50) (Green et al., 2019; Hollingsworth, 2004; Vigod et al., 2018).

Following child removal, parental mental health deteriorated (Wall-Wieler, Roos, Brownell, Nickel, Chateau, and Nixon, 2018d). Compared to women who lost a child to death, child removal was associated with higher rates of maternal anxiety (ARR = 2.51; 95 % CI, 2.40 to 2.63), depression (ARR 1.90; 95 % CI, 1.82 to 1.98), physician contacts for mental health (ARR = 3.01; 95 % CI, 2.91 to 3.12) and psychotropic medication use (ARR = 3.01; 95 % CI, 2.91 to 3.12) (Wall-Wieler, Roos, Bolton, Brownell, Nickel, and Chateau, 2018b). Cross-sectional analysis of health records illustrated birth mothers had higher rates of death by suicide compared to their biological sisters and other mothers in receipt of CPS (RR = 4.46, 95 % CI 1.39–14.33 and RR = 3.45, 95 % CI 1.61–7.40, respectively) (Wall-Wieler, Roos, Brownell, Nickel, Chateau, and Singal, 2018e). Birth mothers were at greatest risk of suicide if they had also been removed from their parents as children (HR = 5.52; 95 % CI 2.91–10.46) (Wall-Wieler et al., 2018a, b, c, d, e, f, g).

Birth parents qualitative testimonies offered insight into complex histories of trauma and the impact of child removal on mental health (Broadhurst et al., 2017; Broadhurst & Mason, 2020; Honey et al., 2019; Memarnia et al., 2015). Birth mothers and fathers spoke of the abandonment of public services and challenges in advocating for appropriate support (Broadhurst et al., 2017) (Philip et al., 2021). The role of being a parent, even following removal, provided meaning and hope for both mothers and fathers (Philip et al., 2021). Support networks were harnessed to renegotiate parental identity, (Hollingsworth, 2004; Sands et al., 2004), although these were less available for birth fathers (Philip et al., 2021).

3.2.2. Physical health ($n = 7$)

Included studies evidenced poor physical health among birth parents before care proceedings, with outcomes worsening after child removal. Compared to mothers without care proceeding involvement, birth mothers' health records reported higher rates of smoking (60 % vs 24 %) and unhealthy weight measurements - clinically underweight (6.9 % vs 3 %) and morbidly obese (5.9 % vs 4.6 %) (Griffiths et al., 2020a, b). Mental health records in the UK found birth mothers had a 2.15 greater risk of death compared to other women accessing similar services (Pearson et al., 2021). Interviews with birth fathers raised multiple long-standing physical conditions, such as chronic pain, asthma, epilepsy, and major dental needs (Philip et al., 2021). In court files, physical health concerns were not cited as reasons for removal (Broadhurst et al., 2017), however mothers' non-attendance at their GP appointments increased the risk of child removal (OR 2.42, 95 % CI 1.42–4.14) (Simkiss et al., 2012).

Following child removal, birth mothers were more likely to self-report their health as 'poor' (OR 1.50, 95 % CI 1.04–2.16) (Kenny et al., 2019). In national mortality data, birth mothers were shown to be at increased risk of dying from amenable (HR 3.04, 95 % CI 2.03–4.57) and preventable causes (HR 3.09, 95 % CI 2.24–4.26) (Wall-Wieler et al., 2018a, b, c, d, e, f, g), including cancer (HR 1.65, 95 % CI 0.72–3.81) and cerebrovascular diseases (HR 1.75, 95 % CI 0.45 to 6.86). Both parents were at increased risk of dying in transport accidents (HR 2.16, 95 % CI 0.26–17.84) and to heart diseases (HR 5.25, 95 % CI 1.08 to 25.43).

3.2.3. Substance misuse and addictions ($n = 16$)

Substances described included methamphetamines, marijuana, alcohol, heroin, and opioid-based prescription medications. Professionals cited concerns of parenting capacity due to substance misuse in care proceeding files (Berger et al., 2010; Broadhurst et al., 2017; Henry et al., 2018) and administrative data reported maternal substance misuse as a risk factor for child removal (OR 8.94; 95 % CI = 5.08–15.71) (Minnes et al., 2008; O'Connor et al., 2020; Sarkola et al., 2007). Mothers accessing specialist drug clinics had high rates of infant entry into care (32–42 % of service users) (Eiden et al., 2007; Sarkola et al., 2007; Wobie et al., 2004) and in ecological analyses, increase opioid prescription rates was associated with more (32 %) children being removed (Quast, 2018). Birth mothers were most at risk if they were living with another drug user (OR 2.71, 95 % CI 1.30–5.56) or had co-morbid mental health challenges (OR 3.9, 95 % CI 1.78–8.55) (Gilchrist & Taylor, 2009).

Birth mothers described using substances as a form of self-medication following child removal, reflecting on the lack of professional support after care proceedings and the function of substances to escape the pain. In some instances, this led to the uptake and increased use of opioids, such as heroin (Broadhurst et al., 2017; Broadhurst & Mason, 2020). Birth fathers described using drugs and alcohol as a form of self-medication, including for pain management, even prior to CPS involvement. It was common for both parents to use drugs, a co-dependency which could produce unhealthy or unsafe environments (Philip et al., 2021).

3.2.4. Intellectual and developmental disabilities ($n = 14$)

Intellectual and developmental disabilities (IDDs) included learning, cognitive and behaviour difficulties, attention deficit hyperactivity disorder (ADHD) and autism spectrum disorders (ASD). Parents with IDD were overrepresented in care proceedings across all included countries (Booth et al., 2005; Booth & Booth, 2004a, 2004b, 2005; Brown et al., 2018; Mayes & Llewellyn, 2012; Rebbe et al., 2020; Welbourne et al., 2017). IDD were frequently mentioned in court files as a concern for parenting capacity (Broadhurst et al., 2017), yet there was lack of clarity around their impact on parenting ability or child welfare (Rice et al., 2021; Sigurjónsdóttir & Rice, 2017, 2018; Tøssebro et al., 2017). A diagnosis of an IDD was often used as a proxy for parenting incapacity, with discriminatory evidence used in court [e.g., parental IQ scores] (Callow et al., 2017). In qualitative interviews, legal professionals disclosed feeling ill-informed for how to support these families and felt training was either absent or insufficient (Cox et al., 2015; Kollinsky et al., 2013). Birth mothers expressed confusion throughout care proceedings with unfamiliar settings and jargon. They described 'mothering differently' and felt powerless within the family justice system (Mayes & Llewellyn, 2012). Birth fathers with IDDs reported challenges articulating emotions, a factor contributing towards professionals' concern (Philip et al., 2021).

In Australia, parents with IDDs had higher rates of keeping their children at home compared to an English sample (59 % vs 10 %) (Booth et al., 2005). In Canada, one in 20 babies born to women with IDDs were discharged into care, a rate 32 times higher than the general population (Brown et al., 2018). Women with comorbid mental illnesses (OR 2.58, 95 % CI 1.90–3.50) and inadequate prenatal care (OR 1.76, 95 % CI 1.32–2.34) were most at risk of removal (Booth and Booth, 2004a, b; Brown et al., 2018; Tøssebro et al., 2017). Court file reviews indicated Indigenous women with IDDs experience particular prejudices and had higher rates of out-of-home placement (60 % vs 48 %) (Collings, Dew, Gordon, Spencer, and Dowse, 2018a).

3.2.5. Reproductive health ($n = 5$)

Birth mothers with children subject to care proceedings were reported to enter motherhood earlier than other women (Brown et al., 2018; Griffiths, Johnson, Broadhurst, Cusworth, Bedston, Jones, Akbari, Lee, Alrouth, Doeblar, John, and Ford, 2020a; Wall-Wieler, Roos, Brownell, Nickel, and Chateau, 2018c). Among women who had infants subject to care proceedings, one in five did not book an antenatal appointment until after 16 weeks (Griffiths, Johnson, Broadhurst, Bedston, Cusworth, Alrouth, Ford, and John, 2020b). Health records indicated that late or no antenatal visit was predictive of out-of-home child placement in Canada (OR 1.76, 95 % CI 1.32–2.34) and Australia (OR 1.61, 95 % CI 1.37–1.90) (Brown et al., 2018; Green et al., 2019).

Findings from UK GP data evidenced maternal use of primary care contraception services was negatively associated with child entry into care (OR 0.64, 95 % CI 0.43–0.97) (Simkiss et al., 2012). Interviews with birth mothers in the UK described grief and loss following child removal as complicating factors in decision-making processes, with women often describing subsequent pregnancies as unplanned (Broadhurst et al., 2017). Less is known about birth fathers' reproductive decision-making choices, although interview data illustrated a desire for more children following removal, comparable to maternal literature (Philip et al., 2021). Repeat pregnancies

Table 2
Supporting health interventions and their evidence base.

Intervention type	Country	Name (if known)	Health focus	Inclusive of fathers (% of sample)	Evidence base	
					Qualitative	Quantitative
Alternative courts	UK	Family Drug and Alcohol Court	Substance misuse	✓ (26 %)	Interviews with 42 parents, 154 court observations and 89 cases.	Randomized trial comparing 90 families in receipt of FDAC. Evaluation of over 400 families in receipt of FDAC vs families in usual court, included self-reported outcomes.
	USA	Family Treatment Court	Substance misuse	✓ (14 %)		
	USA	Mandated treatment	Substance misuse	✓ (12 %)		
	USA	Engaging Moms	Substance misuse	No	Retrospective court file analysis of 477 birth parents.	Randomized pilot study of 62 mothers in the programme.
	UK	Tribunal hearings	Developmental	✓ (4 %)		
Wrap-around services	UK	Multi-site, <i>Positive Choices</i> , <i>MPower</i> , <i>Family Action</i> , <i>Hummingbirds</i>	Flexible	✓ (9 %)	Interviews with 14 birth parents and 5 practitioners.	Self-reported and clinical data from 82 parents.
	USA	Rehabilitation and parenting support	Mental health	✓ (14 %)	Retrospective case file abstraction on 104 mothers accessing rehabilitation and support services.	
	UK	Breaking the Cycle	Flexible		Interviews with 13 birth mothers and 2 social workers.	Self-assessment questionnaires from 25 mothers. Referral forms, progress notes and service use records of 166 women receiving Breaking the Cycle. Self-recorded outcome measures of 9 birth parents.
	Canada	Breaking the Cycle	Substance misuse			
	UK	Reflect	Flexible	✓ (25 %)	Interviews with 4 staff and 16 parents, and analysis of 30 case files.	
	UK	Pause	Flexible		Interviews with 61 women who received Pause.	Secondary analysis of monitoring data capturing 517 women. Questionnaires and programme outcome data reported on 125 mothers.
	Canada	Multi-site <i>Breaking the Cycle</i> , <i>Kids First</i> , <i>Mothering Project</i> , <i>Raising Hope</i> , <i>HerWay Home</i> , <i>Sheway</i> , <i>Maxxine Wright Place</i> , <i>H.E.R</i>	Substance misuse		Interviews with 125 birth mothers, 61 staff and 42 service partners.	
	UK	Hummingbirds	Flexible		Focus groups and interviews with 20 mothers.	
	UK	Option 2	Substance misuse	✓ (13 %)	Interviews with 26 families who received Option 2.	Self-reported outcomes for 31 parents receiving Option 2. Clinical outcome measures on 1408 families receiving FBR.
	USA	Family-based recovery	Substance misuse	✓ (13 %)		
Specialist advocacy/peer support	Australia	Specialist advocacy	Developmental	✓ (30 %)	Structured interviews with 10 birth parents.	Structured interviews with 26 birth mothers. Structured interviews with 14 birth parents. Case study with one practitioner perspective. Interviews with 4 parents and 4 advocates.
	Australia	Peer-support	Developmental		Structured interviews with 26 birth mothers.	
	UK	Specialist advocacy	Developmental	✓ (Not known)	Structured interviews with 14 birth parents.	
	USA	Specialist advocacy	Mental health		Case study with one practitioner perspective.	
	UK	Specialist advocacy	Developmental		Interviews with 4 parents and 4 advocates.	

within quick succession were reflected on by birth mothers as a response to fill an emotional gap and reclaim their motherhood status (Broadhurst et al., 2017). Women described fear of pre-birth assessments and CPS involvement throughout subsequent pregnancies (Mason & Wilkinson, 2021) and were more likely to have inadequate prenatal care (OR 4.29, 95 % CI 3.68 to 5.01) (Wall-Wieler et al., 2019).

3.3. Health interventions for birth parents ($n = 20$)

We grouped health interventions for parents into 3 similar approaches: i) alternative family courts ($n = 4$), ii) wrap-around services ($n = 10$) and iii) specialist advocacy/peer support ($n = 5$). All included interventions were targeted at birth mothers ($n = 20$), with some formally or informally also including birth fathers ($n = 8$). Summary of interventions is in Table 2.

3.3.1. Alternative courts ($n = 4$)

Alternative courts provided different ways of conducting care proceedings, accounting for additional health needs of families before child removal. Most ($n = 3$) addressed parental substance misuse and one evaluated the experiences of parents with IDD. These courts (e.g., Family Drug and Alcohol Court and Engaging Moms Programme) offered therapeutic approaches to proceedings, with multi-disciplinary teams supporting parents through frequent assessments and interventions (Harwin et al., 2018). Parents had regular drug tests and were seen by the same judge throughout. Findings from effectiveness trials demonstrated a reduction in child removal rates and improvement in parental drug misuse (i.e., access to treatment and cessation) compared to families in regular courts (Green et al., 2007; Harwin et al., 2018) (Dakof et al., 2010). Changes were sustained at a 5 year follow up with families (Harwin et al., 2016), although mandated drug treatment for mothers was not shown to reduce the likelihood of returns to court (Rittner & Dozier, 2000). The Scottish Tribunal Hearing system was described as an alternative means of conducting care proceedings (McGhee & Hunter, 2011). The process integrated decision making for children who offend and those in need of care and protection. Unlike care proceedings, hearings consisted of citizen volunteers, the child, birth parents and a social worker. In interviews with parents with IDD, participants reported hearings as less 'scary' and felt positively towards the process's informality. No quantitative evaluation was reported.

3.3.2. Wrap around services ($n = 10$)

Wrap around services provided holistic support for birth parents with care proceeding involvement. These were implemented during high-risk pregnancies (Rutman et al., 2020), as a response to safeguarding concerns (Hanson et al., 2019), and following child removal (Bellew & Peeran, 2017; Cox et al., 2017; McCracken et al., 2020; Roberts et al., 2018). Most services supported mothers ($n = 5$), were delivered via mothers (e.g., in maternity services) and focused on mothers' needs (Andrews et al., 2018). While some supported both parents ($n = 5$), none worked with men in isolation (Roberts et al., 2018). Wrap around services were trauma-informed, relationship-based, and had flexible approaches to outreach and delivery. Services delivered intensive packages of care to support emotional, psychological, and physical needs for between 12 and 24 months (Rutman et al., 2020). Prenatal interventions mitigated some of the negative effects of maternal drug use on mother and baby health outcomes using harm reduction approaches (Rutman et al., 2020). Specialist support for grief, loss, and trauma related to child loss following child removal was also described (McCracken et al., 2020). Multi-agency working with local partnerships encouraged parental engagement with allied healthcare professionals. In Canada, Indigenous liaison workers ensured services were culturally sensitive, exploring the impact of systemic inequities on health service access (Rutman et al., 2020). In UK based services, women were also encouraged to (re)register with GPs to access primary care support (Cox et al., 2017). In the 'Pause' intervention, long-acting reversible contraception was mandated throughout programme (Bellew & Peeran, 2017; Cox et al., 2017; Roberts et al., 2018).

Mixed-method evaluations evidenced improvement in psychological functioning, wellbeing, and relationship capacity for birth parents (Andrews et al., 2018; Bellew & Peeran, 2017; Cook et al., 2014; Cox et al., 2017; Forrester et al., 2016; Hanson et al., 2019; Lewis-Brooke et al., 2017; Mason & Wilkinson, 2021; McCracken et al., 2020; Roberts et al., 2018; Rutman et al., 2020). Interviews with birth mothers reported positive life changes, including healthier living. Women described non-judgmental approaches as key for building trust with services (Forrester et al., 2016; Lewis-Brooke et al., 2017). Both mandating LARCs (McCracken et al., 2020) and/or receiving sexual health advice (Cook et al., 2014; Cox et al., 2017; Roberts et al., 2018) reduced rates of rapid repeat pregnancies for mothers. Findings suggest involving birth fathers would improve whole-family health outcomes (Roberts et al., 2018).

3.3.3. Specialist advocacy/peer support ($n = 5$)

These interventions focused on one-to-one support for parents by individuals with specialist knowledge or experience throughout care proceedings. Most included studies described interventions for parents with IDD ($n = 4$), with one supporting parents with serious mental illnesses (Atkin & Kroese, 2021; Collings, Spencer, Dew, and Dowse, 2018b; Tarleton, 2008; Walton, 2002). Advocacy was based on principles of empowerment, ensuring parents were aware of their rights and supported to exercise them (Collings, Spencer, et al., 2018). Advocates liaised with CPS, court, hospital, and other professionals with birth parents and attended care proceedings. Advocates were a mix of trained volunteers and practitioners with specialist clinical knowledge (Walton, 2002). Birth parents described better understanding of court proceedings with an advocate, yet the lack of structural support for these roles perpetuated feelings powerlessness for parents (Collings, Spencer, Dew, and Dowse, 2018b; Tarleton, 2008).

A peer support model was described for parents with IDD who had experienced domestic violence (Collings et al., 2020), aiming to cultivate emotional and practical support between parents with shared experiences. Narrative interviews with birth mothers involved in peer-support described feelings of comfort and support but highlighted challenges with feasibility, given that all women involved

were facing their own challenges (Collings et al., 2020). We did not find any quantitative evaluations of specialist advocacy or peer support interventions.

4. Discussion

To our knowledge, this is the first study to review parental health in the context of public family care proceedings. Findings describe the known health needs of birth parents; characterise interventions targeted to support these needs and synthesise their evidence base.

4.1. Findings in context

4.1.1. What are the health needs of birth parents?

The included studies illustrate health inequities across all aspects of birth parents' health compared to parents whose children are not subject to care proceedings. Substance misuse and mental health challenges were commonly recorded and considered risk factors for out-of-home child placement (Honey et al., 2019; Sarkola et al., 2007). As evidenced in the qualitative studies, substance use was often a form of self-medication for birth parents experiencing co-morbid conditions and social disadvantages (Broadhurst et al., 2017; Philip et al., 2021). Support services should consider the function of parental substance use, including pain relief for untreated chronic physical health conditions (Canfield et al., 2017). Previous literature has reported associations between poor physical and mental health (Onyeka et al., 2019), but it was not possible to interpret comorbidity with data included in this review. The lack of insight into parental physical health could be due to these needs being unseen by services, and therefore not captured in data, or outcomes of interest being biased towards parental health issues which are a concern in the context of child protection (i.e., health behaviors which pose a 'risk' to a child).

SMI was a consistent risk factor for child removal, particularly for birth mothers (Green et al., 2019; Whitten et al., 2021). In interviews, parents described lifelong health challenges and early childhood adversities (Broadhurst et al., 2017; Memarnia et al., 2015; Philip et al., 2021). There is increasing awareness of the impact early adversity can have on lifelong mental health. Some SMIs, e.g., personality disorders, might be better treated as trauma-based conditions (Bozzatello et al., 2021). Parents with IDD were also overrepresented in care proceedings, with high risk of newborn removal (Booth et al., 2005; Brown et al., 2018). People with SMIs and IDDs have the right to become parents (Broadhurst, Shaw, Kershaw, Harwin, Alrouh, Mason, and Pilling, 2015b). The prompt and high rate of child removal is a health equity issue and brings into focus mainstream service gaps. Unmet health need must not be a key factor in child removal decisions (Broadhurst, Shaw, Kershaw, Harwin, Alrouh, Mason, and Pilling, 2015b).

Child removal triggered health deterioration (Broadhurst & Mason, 2020), including increased rates of anxiety, depression, hospitalisations, and deaths (Wall-Wieler et al., 2018a, b, c, d, e, f, g). Both parents spoke of feeling abandoned by services, with insufficient or no support following proceedings, impacting on their health (Broadhurst & Mason, 2020) (Broadhurst et al., 2017). This review generates clear justification for intensive, specialist support at this time to address factors that led to CPS involvement, and the compounded impact of child removal (Family Rights Group, 2018). There is a notable dearth of information available on birth fathers. As described in qualitative interviews, parental health need can often be interrelated and services must consider 'whole-family' (Woodman et al., 2020) and 'father-specific' need (Philip et al., 2021). Greater involvement of birth fathers should include better recording of paternal status in health data (Lut et al., 2022).

4.1.2. What are the interventions implemented to support these needs?

Included health interventions were based on principles of being relationship-based, trauma-informed, multidisciplinary, family-focused, and longer-term (i.e., up to 2 years) (Cox et al., 2017). Notably, most interventions described in this review were implemented outside of healthcare services and commissioned by local authorities, charities, or short-term innovation funding. This results in high variation between what is available to parents living in different areas (a "postcode lottery") (Mason & Wilkinson, 2021; Rutman et al., 2020). A step forward might be adequate and dedicated funding for services to support birth parents, across health, social and legal services boundaries (Family Rights Group, 2018).

Relationship-based practice with birth parents was important for addressing longstanding distrust of services and improving acceptability and effectiveness of interventions (McCracken et al., 2020; Roberts et al., 2018; Rutman et al., 2020). In the short-term, trusted advocacy or peer support throughout care proceedings helped parents navigate complex court systems and uphold their rights (Collings et al., 2020). Yet advocates supporting parents with IDDs or SMIs faced challenges in the lack of formal structural support for their roles, generating feelings of powerlessness for both parent and advocate (Atkin & Kroese, 2021; Collings, Spencer, Dew, and Dowse, 2018b). Caseworker advocacy roles have been formally implemented to support women who experience intimate partner violence (Rivas et al., 2015), have been trafficked (Westwood et al., 2016), or who are refugees or asylum seekers (Refugee Council, 2022). These examples focus on models of upskilling professionals to respond effectively to specific needs and could be modelled in the context of care proceeding support (Family Rights Group, 2018).

Longer-term wrap-around services provided a way of working with families before, during and after proceedings. These programmes were characterised by offering tailored support via a caseworker who delivered or signposted appropriate services for women and families (Cox et al., 2017; McCracken et al., 2020). The timing of access varied across interventions, with some offering support pre-birth (Rutman et al., 2020; Salford City Council, 2018) and others targeting women following child removal (Cox et al., 2017; McCracken et al., 2020). Services improved birth mothers' emotional wellbeing, relationship capacity and self-esteem (Forrester et al., 2016; Hanson et al., 2019). The rate of rapid repeat pregnancies decreased for women accessing wrap-around support. This was evident in Pause, where LARCs were a condition to the programme (McCracken et al., 2020) and in other services which offered sexual

health education and access (Roberts et al., 2018). As illustrated in a review of parenting interventions for people with SMIs, whole family approaches and family-based work forms crucial components of effective interventions (Radley et al., 2022). The UK ‘Supporting Families’ is an example of such whole-family caseworker model (UK Gov, 2022).

FDAC evaluations evidenced the benefit of multi-disciplinary support for parents who misuse substances (Harwin et al., 2018). Results demonstrated a reduction in parental substance misuse and improvement in family reunification rates (Harwin et al., 2016). Whilst these findings are promising, there are known challenges with non-linear recovery from drug and alcohol addictions (Laudet et al., 2002). The alternative courts described in this review relied on parental abstinence, rather than harm-reduction approaches implemented in other contexts (Boyd et al., 2022). Harm-reduction approaches to maternity care for drug using pregnant woman in this review illustrated promising health outcomes for both mother and child (Rutman et al., 2020). Providing a safe environment for wrap-around prenatal care mitigated many of the negative effects of maternal drug use and improved service engagement for families (Rutman et al., 2020). The antenatal period offers an opportunity for targeted intervention, yet we know that birth parents may be reluctant to engage with services due to fear of child removal (Broadhurst et al., 2017). More research is needed to understand the role of maternity services in supporting birth parents at risk of child removal (Griffiths, Johnson, Broadhurst, Bedston, Cusworth, Alrouh, Ford, and John, 2020b) (see Table 3).

4.2. Limitations

We were unable to draw strong conclusions on the effectiveness of included interventions and further systematic review methods and meta-analysis is needed. We did not include research on social need, such as experiences of homelessness, poverty, and violence. These findings would undoubtedly be relevant to the health outcomes of birth parents and should be explored further. This review excluded publications not available in English, which might have limited the international significance of findings. As we only included articles describing child placement, results may also underestimate the role of CPS in supporting parental health needs for families who do not undergo care proceedings.

4.3. Implications of findings

Findings from this review can be used to inform the development and delivery of services supporting the health needs of birth parents who have children subject to care proceedings. Implications of findings are outlined in Table 3.

4.4. Conclusion

Parents who have children subject to care proceedings have complex health needs that often manifest before CPS involvement. The included studies strongly suggest health issues are exacerbated by child removal, including avoidable mortality. There are models that have been designed, implemented, and tested to support birth parents’ health using relationship-based, trauma-informed, multidisciplinary, family-focused, and long-term approaches.

Table 3
Key implications of scoping review.

Implications for practitioners
• Ask service users about parental status and family planning • Develop trust with families by working in a relationship- and strengths-based way • Implement trauma-informed approaches to working with birth parents • Acknowledge inter-related health need by considering ‘whole-family’ health • Be an advocate for the rights of birth parents • Utilise and strengthen local networks of support
Implications for funders
• Consider commissioning longer-term, holistic support across social care and health • Blueprinting relationship-based, trauma-informed, long-term support for parents at risk of child removal or who have had a child removed. • Investment in preventative (pre-birth) intervention for adults with complex needs • Implement enhanced training programmes for practitioners working with birth parents
Directions for future research
• Greater involvement of birth fathers in research and intervention development • Understand experiences of health from birth parents’ perspective • Develop more individual-level data-linkages between CPS and parents’ healthcare records • Research into the role of maternity services supporting complex health needs

Funding

CG is supported by the Economic and Social Research Council (ESRC) UBEL Doctoral Training Programme (ref: ES/P000592/1). JW, CP, and RL are in part supported by the NIHR Children and Families Policy Research Unit (PR-PRU-1217-21301).

Data availability

No data was used for the research described in the article.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.chiabu.2023.106160>.

References

- Andrews, N. C. Z., Motz, M., Pepler, D. J., Jeong, J. J., & Khoury, J. (2018). Engaging mothers with substance use issues and their children in early intervention: Understanding use of service and outcomes. *Child Abuse & Neglect*, 83, 10–20. <https://doi.org/10.1016/j.chiabu.2018.06.011>
- Atkin, C., & Kroese, B. S. (2021). Exploring the experiences of independent advocates and parents with intellectual disabilities, following their involvement in child protection proceedings. *Disability and Society*, 1–23. <https://doi.org/10.1080/09687599.2021.1881884>
- Barlow, J., Simkiss, D., & Stewart-Brown, S. (2006). Interventions to prevent or ameliorate child physical abuse and neglect: Findings from a systematic review of reviews. *Journal of Children's Services*, 1(3), 6–28. <https://doi.org/10.1108/17466660200600020>
- Bedston, S., Philip, G., Youansamouth, L., Clifton, J., Broadhurst, K., Brandon, M., & Hu, Y. (2019). Linked lives: Gender, family relations and recurrent care proceedings in England. *Children and Youth Services Review*. <https://doi.org/10.1016/j.childyouth.2019.104392>
- Bellew, R., & Peeran, U. (2017). *After adoptions breaking the cycle programme: An evaluation of the two year pilot, September 2014 to August 2016 (Issue September 2017)*. Coram.
- Berger, L. M., Slack, K. S., Waldfogel, J., & Bruch, S. K. (2010). Caseworker-perceived caregiver substance abuse and child protective services outcomes. *Child Maltreatment*, 15(3), 199–210. <https://doi.org/10.1177/1077559510368305>
- Berlin, M., Vinnerljung, B., & Hjern, A. (2011). School performance in primary school and psychosocial problems in young adulthood among care leavers from long term foster care. *Children and Youth Services Review*, 33(12), 2489–2497. <https://doi.org/10.1016/j.childyouth.2011.08.024>
- Booth, T., & Booth, W. (2004). *Parents with learning difficulties child protection and the courts: A report (Issue March, pp. 1–190)*. The Nuffield Foundation. <http://www.supportedparenting.co.uk/projects/courts.html>
- Booth, T., & Booth, W. (2005). Parents with learning difficulties in the child protection system: Experiences and perspectives. *Journal of Intellectual Disabilities*. <https://doi.org/10.1177/1077559505053922>
- Booth, T., Booth, W., & McConnell, D. (2005). The prevalence and outcomes of care proceedings involving parents with learning difficulties in the family courts. *Journal of Applied Research in Intellectual Disabilities*. <https://doi.org/10.1111/j.1468-3148.2004.00204.x>
- Booth, W., & Booth, T. (2004). A family at risk: Multiple perspectives on parenting and child protection. *British Journal of Learning Disabilities*. <https://doi.org/10.1111/j.1468-3156.2004.00263.x>
- Boyd, J., Maher, L., Austin, T., Lavalley, J., Kerr, T., & McNeil, R. (2022). Mothers who use drugs: Closing the gaps in harm reduction response amidst the dual epidemics of overdose and violence in a Canadian urban setting. *American Journal of Public Health*, 112, S191–S198. <https://doi.org/10.2105/AJPH.2022.306776>
- Bozzatello, P., Rocca, P., Baldassarri, L., Bosia, M., & Bellino, S. (2021). The role of trauma in early onset borderline personality disorder: A biopsychosocial perspective. *Frontiers in Psychiatry*, 12, Article 721361. <https://doi.org/10.3389/fpsy.2021.721361>
- Broadhurst, K., Alrouh, B., Yeend, E., Harwin, J., Shaw, M., Pilling, M., Mason, C., & Kershaw, S. (2015). Connecting events in time to identify a hidden population: Birth mothers and their children in recurrent care proceedings in England. *The British Journal of Social Work*, 45(8), 2241–2260.
- Broadhurst, K., & Mason, C. (2013). Maternal outcasts: Raising the profile of women who are vulnerable to successive, compulsory removals of their children—A plea for preventative action. *Journal of Social Welfare and Family Law*. <https://doi.org/10.1080/09649069.2013.805061>
- Broadhurst, K., & Mason, C. (2020). Child removal as the gateway to further adversity: Birth mother accounts of the immediate and enduring collateral consequences of child removal. *Qualitative Social Work*. <https://doi.org/10.1177/1473325019893412>
- Broadhurst, K., Mason, M. C., Bedston, S., Alrouh, B., Morriss, L., McQuarrie, M. T., Palmer, M., Shaw, M., Harwin, J., & Kershaw, M. S. (2017). *Vulnerable birth mothers and recurrent care proceedings*. Lancaster: University of Lancaster.
- Broadhurst, K., Shaw, M., Kershaw, S., Harwin, J., Alrouh, B., Mason, C., & Pilling, M. (2015). Vulnerable birth mothers and repeat losses of infants to public care: Is targeted reproductive health care ethically defensible? *Journal of Social Welfare and Family Law*, 37(1), 84–98. <https://doi.org/10.1080/09649069.2015.998007>
- Brown, H. K., Potvin, L. A., Lunskey, Y., & Vigod, S. N. (2018). Maternal intellectual or developmental disability and newborn discharge to protective services. *Pediatrics*, 142(6). <https://doi.org/10.1542/peds.2018.1416>
- Bywaters, P., Brady, G., Sparks, T., & Bos, E. (2016). Inequalities in child welfare intervention rates: The intersection of deprivation and identity. *Child and Family Social Work*, 21(4), 452–463. <https://doi.org/10.1111/cfs.12161>
- Callow, E., Tahir, M., & Feldman, M. (2017). Judicial reliance on parental IQ in appellate-level child welfare cases involving parents with intellectual and developmental disabilities. *Journal of Applied Research in Intellectual Disabilities*. <https://doi.org/10.1111/jar.12296>
- Canfield, M., Radcliffe, P., Marlow, S., Boreham, M., & Gilchrist, G. (2017). Maternal substance use and child protection: A rapid evidence assessment of factors associated with loss of child care. *Child Abuse and Neglect*, 70, 11–27. <https://doi.org/10.1016/j.chiabu.2017.05.005>
- Collings, S., Dew, A., Gordon, T., Spencer, M., & Dowse, L. (2018). Intersectional disadvantage: Exploring differences between aboriginal and non-aboriginal parents with intellectual disability in the New South Wales child protection system. *Journal of Public Child Welfare*. <https://doi.org/10.1080/15548732.2017.1379456>
- Collings, S., Spencer, M., Dew, A., & Dowse, L. (2018). 'She was there if I needed to talk or to try and get my point across': Specialist advocacy for parents with intellectual disability in the Australian child protection system. *Australian Journal of Human Rights*. <https://doi.org/10.1080/1323238X.2018.1478595>
- Collings, S., Strnadová, I., Loblinz, J., & Danker, J. (2020). Benefits and limits of peer support for mothers with intellectual disability affected by domestic violence and child protection. *Disability and Society*. <https://doi.org/10.1080/09687599.2019.1647150>
- Coman, W., & Devaney, J. (2011). Reflecting on outcomes for looked-after children: An ecological perspective. *Child Care in Practice*, 17(1), 37–53. <https://doi.org/10.1080/13575279.2010.522976>
- Cook, J. A., Steigman, P. J., & Jonikas, J. A. (2014). Outcomes of programs serving mothers with psychiatric disabilities and their young children: A multisite case file abstraction study. *Psychiatric Rehabilitation Journal*, 37(3), 232–241. <https://doi.org/10.1037/prj0000075>
- Courtney, M. E., Dworsky, A. L., & Perez, A. (2007). Midwest evaluation of the adult functioning of former foster youth: Outcomes at age 21. *Social Work*, May, 1–93.
- Cox, P., Barratt, C., Blumenfeld, F., Rahemtulla, Z., Taggart, D., & Turton, J. (2017). Reducing recurrent care proceedings: Initial evidence from new interventions. *Journal of Social Welfare and Family Law*, 39(3), 332–349. <https://doi.org/10.1080/09649069.2017.1345083>

- Cox, R., Stenfort Kroese, B., & Evans, R. (2015). Solicitors' experiences of representing parents with intellectual disabilities in care proceedings: Attitudes, influence and legal processes. *Disability and Society*. <https://doi.org/10.1080/09687599.2015.1005730>
- Dakof, G. A., Cohen, J. B., Henderson, C. E., Duarte, E., Boustani, M., Blackburn, A., Venzer, E., & Hawes, S. (2010). A randomized pilot study of the engaging moms program for family drug court. *Journal of Substance Abuse Treatment*, 38(3), 263–274. <https://doi.org/10.1016/j.jsat.2010.01.002>
- Eiden, R. D., Foote, A., & Schuetz, P. (2007). Maternal cocaine use and caregiving status: Group differences in caregiver and infant risk variables. *Addictive Behaviors*, 32(3), 465–476. <https://doi.org/10.1016/j.addbeh.2006.05.013>
- Family Rights Group. (2018). *Care crisis review: Options for change*.
- Forrester, D., Holland, S., Williams, A., & Copello, A. (2016). Helping families where parents misuse drugs or alcohol? A mixed methods comparative evaluation of an intensive family preservation service. *Child and Family Social Work*, 21(1), 65–75. <https://doi.org/10.1111/cfs.12111>
- Gilbert, N., Parton, N., & Skiveness, M. (2011). Child protection systems: International trends and orientations. In *Oxford University Press*. <https://doi.org/10.1093/acprof:oso/9780199793358.001.0001>
- Gilchrist, G., & Taylor, A. (2009). Drug-using mothers: Factors associated with retaining care of their children. *Drug and Alcohol Review*, 28(2), 175–185. <https://doi.org/10.1111/j.1465-3362.2008.00017.x>
- Grant, C., Radley, J., Lacey, R., & Woodman, J. (2021). Role of public services in supporting the health needs of birth parents whose children are subject to care proceedings: A scoping review. *Open Science Framework*. <https://osf.io/ykq2v/>.
- Green, B. L., Furrer, C., Worcel, S., Burrus, S., & Finigan, M. W. (2007). How effective are family treatment drug courts? Outcomes from a four-site national study. *Child Maltreatment*, 12(1), 43–59. <https://doi.org/10.1177/1077559506296317>
- Green, M. J., Kariuki, M., Chilvers, M., Butler, M., Katz, I., Burke, S., Tzoumakis, S., Laurens, K. R., Harris, F., & Carr, V. J. (2019). Inter-agency indicators of out-of-home-care placement by age 13–14 years: A population record linkage study. *Child Abuse & Neglect*, 93, 91–102. <https://doi.org/10.1016/j.chiabu.2019.04.013>
- Griffiths, L., Johnson, R., Broadhurst, K., Cusworth, L., Bedston, S., Jones, K., Akbari, A., Lee, A., Alrouth, B., Doeblen, S., John, A., & Ford, D. (2020). *Born into care: One thousand mothers in care proceedings in Wales: Maternal health, wellbeing and pregnancy outcomes*. June.
- Griffiths, L., Johnson, R. D., Broadhurst, K., Bedston, S., Cusworth, L., Alrouth, B., Ford, D. V., & John, A. (2020). Maternal health, pregnancy and birth outcomes for women involved in care proceedings in Wales: A linked data study. *BMC Pregnancy and Childbirth*, 20(1). <https://doi.org/10.1186/s12884-020-03370-4>
- Hanson, K. E., Duryea, E. R., Painter, M., Vanderploeg, J. J., & Saul, D. H. (2019). Family-based recovery: An innovative collaboration between community mental health agencies and child protective services to treat families impacted by parental substance use. *Child Abuse Review*, 28(1), 69–81. <https://doi.org/10.1002/car.2545>
- Harwin, J., Alrouth, B., Ryan, M., McQuarrie, T., Golding, L., Broadhurst, K., Tunnard, J., & Swift, S. (2016). After FDAC: Outcomes 5 years later, final report (Issue December). <http://wp.lancs.ac.uk/cj-fdac/publications/>.
- Harwin, J., Ryan, M., & Broadhurst, K. (2018). How does FDAC succeed with parents with substance misuse Problems? Exploring relational practices within the english family drug and alcohol court. *Child Abuse Review*, 27(4), 266–279. <https://doi.org/10.1002/car.2521>
- Henry, C., Liner-Jigamian, N., Carnochan, S., Taylor, S., & Austin, M. J. (2018). Parental substance use: How child welfare workers make the case for court intervention. *Children and Youth Services Review*, 93, 69–78. <https://doi.org/10.1016/j.childyouth.2018.07.003>
- Hollingsworth, L. D. (2004). Persistent severe mental illness. *Social Work Research*, 28(4), 199–209.
- Honey, A., Miceli, M., & Mayes, R. (2019). Living with mental illness and child removal. *Advances in Mental Health*. <https://doi.org/10.1080/18387357.2019.1614884>
- Källquist, A., & Salzmann-Erikson, M. (2019). Experiences of having a parent with serious mental illness: An interpretive meta-synthesis of qualitative literature. *Journal of Child and Family Studies*, 28(8), 2056–2068. <https://doi.org/10.1007/s10826-019-01438-0>
- Kenny, K. S., Ranville, F., Green, S. L., Duff, P., Braschel, M., Abrahams, R., & Shannon, K. (2019). Family separation and maternal self-rated health: Evidence from a prospective cohort of marginalized mothers in a Canadian setting. *Maternal and Child Health Journal*, 23(9), 1232–1239. <https://doi.org/10.1007/s10995-019-02762-z>
- Kohl, P. L., Jonson-Reid, M., & Drake, B. (2011). Maternal mental illness and the safety and stability of maltreated children. *Child Abuse and Neglect*, 35(5). <https://doi.org/10.1016/j.chiabu.2011.01.006>
- Kollinsky, L., Simonds, L. M., & Nixon, J. (2013). A qualitative exploration of the views and experiences of family court magistrates making decisions in care proceedings involving parents with learning disabilities. *British Journal of Learning Disabilities*. <https://doi.org/10.1111/j.1468-3156.2012.00726.x>
- Kratky, N., & Schröder-Abé, M. (2018). How are parental functioning and single parenthood associated with court outcomes? An analysis of child protection cases. *Child Abuse and Neglect*, 84, 95–105. <https://doi.org/10.1016/j.chiabu.2018.07.015>
- Laudet, A. B., Savage, R., & Mahmood, D. (2002). Pathways to long-term recovery: A preliminary investigation. *Journal of Psychoactive Drugs*, 34(3), 305–311.
- Lewis-Brooke, S., Bell, L., Herring, R., Lehan, O'Farrell-Pearce, Quinn, & So. (2017). Mothers apart: An action research project based on partnership between a local authority and a university in London, England. *Revista de Asistență Socială*, XVI, 1–11.
- Lut, I., Harron, K., Hardelid, P., O'Brien, M., & Woodman, J. (2022). 'What about the dads?' Linking fathers and children in administrative data: A systematic scoping review. *Big Data and Society*, 9(1). <https://doi.org/10.1177/20539517211069299>
- Mason, C., & Wilkinson, J. (2021). Services for parents who have experienced recurrent care proceedings: Where are we now?. In *Findings from mapping of locally developed services in England*. Dartington: Research in Practice.
- Mayes, R., & Llewellyn, G. (2012). Mothering differently: Narratives of mothers with intellectual disability whose children have been compulsorily removed. *Journal of Intellectual and Developmental Disability*. <https://doi.org/10.3109/13668250.2012.673574>
- McCracken, K., Priest, S., Fitzsimons, A., Bracewell, K., Torchia, K., Parry, W., & Stanley, N. (2020). *Evaluation of pause (issue July)*.
- McGhee, J., & Hunter, S. (2011). The Scottish children's hearings tribunals system: A better forum for parents with learning disabilities? *Journal of Social Welfare and Family Law*. <https://doi.org/10.1080/09649069.2011.626245>
- Memarnia, N., Nolte, L., Norris, C., & Harborne, A. (2015). 'It felt like it was night all the time': Listening to the experiences of birth mothers whose children have been taken into care or adopted. *Adoption and Fostering*, 39(4), 303–317. <https://doi.org/10.1177/0308575915611516>
- Minnes, S., Singer, L. T., Humphrey-Wall, R., & Satayathum, S. (2008). Psychosocial and behavioral factors related to the post-partum placements of infants born to cocaine-using women. *Child Abuse & Neglect*, 32(3), 353–366. <https://doi.org/10.1016/j.chiabu.2007.12.002>
- Munn, Z., Peters, M. D. J., Stern, C., Tufanaru, C., McArthur, A., & Aromataris, E. (2018). Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Medical Research Methodology*. <https://doi.org/10.1186/s12874-018-0611-x>
- Munro, E. R., & Ward, H. (2008). Balancing parents' and very young children's rights in care proceedings: Decision-making in the context of the human rights act 1998. *Child and Family Social Work*, 13(2), 227–234. <https://doi.org/10.1111/j.1365-2206.2007.00542.x>
- Nevriana, A., Pierce, M., Dalman, C., Wicks, S., Hasselberg, M., Hope, H., Abel, K. M., & Kosidou, K. (2020). Association between maternal and paternal mental illness and risk of injuries in children and adolescents: Nationwide register based cohort study in Sweden. *BMJ (Clinical Research Ed.)*, 369, Article m853. <https://doi.org/10.1136/bmj.m853>
- O'Connor, A., Harris, E., Seeber, C., Hamilton, D., Fisher, C., & Sachmann, M. (2020). Methamphetamine use in pregnancy, child protection, and removal of infants: Tertiary Centre experience from Western Australia. *Midwifery*, 83, Article 102641. <https://doi.org/10.1016/j.midw.2020.102641>
- Onyeka, I. N., Collier Hoegh, M., Nâheim Eien, E. M., Nwaru, B. I., & Melle, I. (2019). Comorbidity of physical disorders among patients with severe mental illness with and without substance use disorders: A systematic review and meta-analysis. *Journal of Dual Diagnosis*, 15(3), 192–206. <https://doi.org/10.1080/15504263.2019.1619007>
- Park, J. M., Solomon, P., & Mandell, D. S. (2006). Involvement in the child welfare system among mothers with serious mental illness. *Psychiatric Services (Washington, D.C.)*, 57(4), 493–497. <https://doi.org/10.1176/ps.2006.57.4.493>
- Pearson, R., Grant, C., Wijlaars, L., Finch, E., Bedston, S., Broadhurst, K., & Gilbert, R. (2021). Mental health service use among mothers involved in public family law proceedings: Linked data cohort study in South London 2007–2019. *Social Psychiatry and Psychiatric Epidemiology*, Article 0123456789. <https://doi.org/10.1007/s00127-022-02221-1>

- Pearson, R., Jewell, A., Wijlaars, L., Bedston, S., Finch, E., Bradley, K., Downs, J., & Gilbert, R. (2020). Linking data on women in public family law court proceedings concerning their children to mental health service records in South London. *International Journal of Population Data Science*, (August).
- Pearson, R. J., Jay, M. A., Wijlaars, L. P. M. M., De Stavola, B., Syed, S., Bedston, S. J., & Gilbert, R. (2020). Association between health indicators of maternal adversity and the rate of infant entry to local authority care in England: A longitudinal ecological study. *BMJ Open*. <https://doi.org/10.1136/bmjopen-2019-036564>
- Philip, G., Bedston, S., Youansamouth, L., Clifton, J., Broadhurst, K., Brandon, M., Hu, Y., Author, C., & Philip, G. (2021). 'UP AGAINST IT': *Understanding fathers' repeat appearance in local authority care*. The Nuffield Foundation.
- Pierce, M., Hope, H. F., Kolade, A., Gellatly, J., Osam, C. S., Perchard, R., Kosidou, K., Dalman, C., Morgan, V., Di Prinzio, P., & Abel, K. M. (2020). Effects of parental mental illness on children's physical health: Systematic review and meta-analysis. *British Journal of Psychiatry*, 217(1), 354–363. <https://doi.org/10.1192/bjp.2019.216>
- Public Health England. (2014). Health matters: Public health issues. <https://www.gov.uk/government/collections/health-matters-public-health-issues>.
- Quast, T. (2018). State-level variation in the relationship between child removals and opioid prescriptions. *Child Abuse & Neglect*, 86, 306–313. <https://doi.org/10.1016/j.chiabu.2018.10.001>
- Radley, J., Sivarajah, N., Moltrecht, B., Klampe, M. L., Hudson, F., Delahay, R., Barlow, J., & Johns, L. C. (2022). A scoping review of interventions designed to support parents with mental illness that would be appropriate for parents with psychosis. *Frontiers in Psychiatry*, 12(January). <https://doi.org/10.3389/fpsy.2021.787166>
- Rebbe, R., Brown, S. E., Matter, R. A., & Mienko, J. A. (2020). Prevalence of births and interactions with child protective services of children born to mothers diagnosed with an intellectual and/or developmental disability. *Maternal and Child Health Journal*. <https://doi.org/10.1007/s10995-020-03105-z>
- Refugee Council. (2022). *Helping refugees and asylum seekers to find healthcare*. Refugee Council. <https://www.refugeecouncil.org.uk/projects/helping-refugees-and-asylum-seekers-to-find-healthcare/>.
- Rice, J. G., Bjargardóttir, H. B., & Sigurjónsdóttir, H. B. (2021). Child protection, disability and obstetric violence: Three case studies from Iceland. *International Journal of Environmental Research and Public Health*. <https://doi.org/10.3390/ijerph18010158>
- Rittner, B., & Dozier, C. D. (2000). Effects of court-ordered substance abuse treatment in child protective services cases. *Social Work*, 45(2), 131–140. <https://doi.org/10.1093/sw/45.2.131>
- Rivas, C., Ramsay, J., Sadowski, L., Davidson, L. L., Dunne, D., Eldridge, S., Hegarty, K., Taft, A., & Feder, G. (2015). Advocacy interventions to reduce or eliminate violence and promote the physical and psychosocial well-being of women who experience intimate partner abuse. *Cochrane Database of Systematic Reviews*, 12. <https://doi.org/10.1002/14651858.cd005043.pub3>
- Roberts, L., Maxwell, N., Messenger, M. R., & Palmer, M. C. (2018). *Evaluation of reflect in Gwent final report (Issue September)*.
- Rutman, D., Hubberstey, C., Poole, N., Schmidt, R. A., & Van Bibber, M. (2020). Multi-service prevention programs for pregnant and parenting women with substance use and multiple vulnerabilities: Program structure and clients' perspectives on wraparound programming. *BMC Pregnancy and Childbirth*, 20(1). <https://doi.org/10.1186/s12884-020-03109-1>
- Salford City Council. (2018). *Salford strengthening families*.
- Salzer, M. S., Berg, K. L., Kaplan, K., & Brusilovskiy, E. (2020). Custody challenges experienced by parents with serious mental illnesses outside of child protective services proceedings. *Psychiatric Rehabilitation Journal*. <https://doi.org/10.1037/prj0000448>
- Sands, R. G., Koppelman, N., & Solomon, P. (2004). Maternal custody status and living arrangements of children of women with severe mental illness. *Health and Social Work*, 29(4), 317–325. <https://doi.org/10.1093/hsw/29.4.317>
- Sarkola, T., Kahila, H., Gissler, M., & Halmesmäki, E. (2007). Risk factors for out-of-home custody child care among families with alcohol and substance abuse problems. *Acta Paediatrica, International Journal of Paediatrics*, 96(11), 1571–1576. <https://doi.org/10.1111/j.1651-2227.2007.00474.x>
- Sheehan, R., & Levine, G. (2005). Parents with mental illness: Decision-making in Australian children's court cases involving parents with mental health problems. *Journal of Social Welfare and Family Law*, 27(1), 17–30. <https://doi.org/10.1080/09649060500085842>
- Sigurjónsdóttir, H. B., & Rice, J. G. (2017). 'Framed': Terminating the parenting rights of parents with intellectual disability in Iceland. *Journal of Applied Research in Intellectual Disabilities*. <https://doi.org/10.1111/jar.12301>
- Sigurjónsdóttir, H. B., & Rice, J. G. (2018). Evidence of neglect as a form of structural violence: Parents with intellectual disabilities and custody deprivation. *Social Inclusion*. <https://doi.org/10.17645/si.v6i2.1344>
- Simkiss, D. E., Spencer, N. J., Stallard, N., & Thorogood, M. (2012). Health service use in families where children enter public care: A nested case control study using the general practice research database. *BMC Health Services Research*, 12(1). <https://doi.org/10.1186/1472-6963-12-65>
- Skinner, G. C. M., Bywaters, P. W. B., Bilson, A., Duschinsky, R., Clements, K., & Hutchinson, D. (2021). The 'toxic trio' (domestic violence, substance misuse and mental ill-health): How good is the evidence base? *Children and Youth Services Review*, 120(August 2020), Article 105678. <https://doi.org/10.1016/j.childyouth.2020.105678>
- Tarleton, B. (2008). Specialist advocacy services for parents with learning disabilities involved in child protection proceedings. *British Journal of Learning Disabilities*. <https://doi.org/10.1111/j.1468-3156.2007.00479.x>
- Tøssebro, J., Midjo, T., Paulsen, V., & Berg, B. (2017). Prevalence, trends and custody among children of parents with intellectual disabilities in Norway. *Journal of Applied Research in Intellectual Disabilities*. <https://doi.org/10.1111/jar.12304>
- Tricco, A. C., Lillie, E., Zarin, W., O'Brien, K. K., Colquhoun, H., Levac, D., Straus, S. E., ... (2018). PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Annals of Internal Medicine*, 169(7), 467–473. <https://doi.org/10.7326/M18-0850>
- UK Gov. (2022). *The national supporting families outcome framework*. GOV.UK. <https://www.gov.uk/government/publications/supporting-families-programme-guidance-2022-to-2025/chapter-3-the-national-supporting-families-outcome-framework>.
- UK Government. (2022). Levelling up white paper. <https://www.gov.uk/government/publications/levelling-up-the-united-kingdom>.
- UK Public General Acts. (1989). Children Act 1989. <https://www.legislation.gov.uk/ukpga/1989/41/contents>.
- United Nations. (1989). *In United nations treaty series. Convention on the rights of the child* (pp. 3–178).
- Vigod, S. N., Laursen, T. M., Ranning, A., Nordentoft, M., & Munk-Olsen, T. (2018). Out-of-home placement to age 18 years in children exposed to a postpartum mental disorder. *Acta Psychiatrica Scandinavica*, 138(1), 35–43. <https://doi.org/10.1111/acps.12890>
- Wall-Wieler, E., Bolton, J., Liu, C., Wilcox, H., Roos, L. L., & Hjern, A. (2018). Intergenerational involvement in out-of-home care and death by suicide in Sweden: A population-based cohort study. *Journal of Affective Disorders*, 238(March), 506–512. <https://doi.org/10.1016/j.jad.2018.06.022>
- Wall-Wieler, E., Kenny, K., Lee, J., Thiessen, K., Morris, M., & Roos, L. L. (2019). Prenatal care among mothers involved with child protection services in Manitoba: A retrospective cohort study. *CMAJ*, 191(8), E209–E215. <https://doi.org/10.1503/cmaj.181002>
- Wall-Wieler, E., Roos, L. L., Bolton, J., Brownell, M., Nickel, N., & Chateau, D. (2018). Maternal mental health after custody loss and death of a child: A retrospective cohort study using linkable administrative data. *Canadian Journal of Psychiatry*, 63(5), 322–328. <https://doi.org/10.1177/0706743717738494>
- Wall-Wieler, E., Roos, L. L., Brownell, M., Nickel, N., Chateau, D., & Singal, D. (2018). Suicide attempts and completions among mothers whose children were taken into care by child protection services: A cohort study using linkable administrative data. *Canadian Journal of Psychiatry*, 63(3), 170–177. <https://doi.org/10.1177/0706743717741058>
- Wall-Wieler, E., Roos, L. L., Brownell, M., Nickel, N. C., & Chateau, D. (2018). Predictors of having a first child taken into care at birth: A population-based retrospective cohort study. *Child Abuse and Neglect*, 76(October 2017), 1–9. <https://doi.org/10.1016/j.chiabu.2017.09.033>
- Wall-Wieler, E., Roos, L. L., Brownell, M., Nickel, N. C., Chateau, D., & Nixon, K. (2018). Postpartum depression and anxiety among mothers whose child was placed in Care of Child Protection Services at birth: A retrospective cohort study using linkable administrative data. *Maternal and Child Health Journal*, 22(10), 1393–1399. <https://doi.org/10.1007/s10995-018-2607-x>
- Wall-Wieler, E., Roos, L. L., Nickel, N. C., Chateau, D., & Brownell, M. (2018). Mortality among mothers whose children were taken into care by child protection services: A discordant sibling analysis. *American Journal of Epidemiology*, 187(6), 1182–1188.
- Wall-Wieler, E., Vinnerljung, B., Liu, C., Roos, L. L., & Hjern, A. (2018). Avoidable mortality among parents whose children were placed in care in Sweden: A population-based study. *Journal of Epidemiology and Community Health*, 72(12), 1091–1098. <https://doi.org/10.1136/jech-2018-210672>

- Walton, M. K. (2002). Advocacy and leadership when parental rights and child welfare collide: The role of the advanced practice nurse. *Journal of Pediatric Nursing*, 17 (1), 49–58. <https://doi.org/10.1053/jpdn.2002.30928>
- Welbourne, P., Macdonald, P., & Bates, P. (2017). *Getting it right in time : Parents who lack litigation capacity in care proceedings (Issue July)*. Nuffield Foundation. <https://www.nuffieldfoundation.org/project/getting-it-right-in-time-parents-who-lack-litigation-capacity-in-public-law>.
- Westwood, J., Howard, L. M., Stanley, N., Zimmerman, C., Gerada, C., & Oram, S. (2016). Access to, and experiences of, healthcare services by trafficked people: Findings from a mixed-methods study in England. *British Journal of General Practice*, 66(652), e794–e801. <https://doi.org/10.3399/bjgp16X687073>
- Whitten, T., Dean, K., Li, R., Laurens, K. R., Harris, F., Carr, V. J., & Green, M. J. (2021). Earlier contact with child protection services among children of parents with criminal convictions and mental disorders. *Child Maltreatment*, 26(1), 63–73. <https://doi.org/10.1177/1077559520935204>
- Wobie, K., Eyler, F. D., Garvan, C. W., Hou, W., & Behnke, M. (2004). Prenatal cocaine exposure: An examination of out-of-home placement during the first year of life. *Journal of Drug Issues*, 34(1), 77–94. <https://doi.org/10.1177/002204260403400104>
- Woodman, J., Simon, A., Hauari, H., & Gilbert, R. (2020). A scoping review of ‘think-family’ approaches in healthcare settings. *Journal of Public Health (United Kingdom)*. <https://doi.org/10.1093/pubmed/fdy210>